

Response of *Syngonium podophyllum* L. 'White Butterfly' shoot cultures to alternative media additives and gelling agents, and flow cytometric analysis of regenerants

JAIME A. TEIXEIRA DA SILVA

¹ Faculty of Agriculture and Graduate School of Agriculture, Kagawa University, Miki-cho, Kagawa, 761-0795, Japan. Tel: +81-87-891-3008, Fax: +81-87-891-3021, email: jaimetex@yahoo.com

² Present address: P. O. Box 7, Miki-cho post office, Ikenobe 3011-2, Kagawa-ken, 761-0799, Japan

Manuscript received: 9 December 2014. Revision accepted: 20 January 2015.

Abstract. Teixeira da Silva JA. 2015. Response of *Syngonium podophyllum* L. 'White Butterfly' shoot cultures to alternative media additives and gelling agents, and flow cytometric analysis of regenerants. *Nusantara Bioscience* 7: 26-32. *Syngonium podophyllum* L. (arrowhead vine) is a popular leafy indoor pot plant whose tissue culture has been established, primarily through *in vitro* shoot culture, but several interesting aspects have not yet been explored. In this study, cv. 'White Butterfly' was used to investigate the response of shoot formation to alternative gelling agents and media additives. Gellan gum (Gelrite[®]) at 2 g/L resulted in greater leaf production, plantlet fresh weight and higher chlorophyll content (SPAD value) than all other gelling agents tested, including agar, Bacto agar, phytagel, oatmeal agar, potato dextrose agar, barley starch and corn starch, when on a basal Hyponex[®] (NPK = 6.5: 6: 19; 3 g/L) medium. Several alternative liquid medium additives tested (low and full fat milk, Coca-Cola[®], coffee, Japanese green, Oolong and Darjeeling teas) negatively impacted plant growth, stunted roots and decreased chlorophyll content (SPAD value) of leaves. Plant growth on medium with refined sucrose or table sugar responded similarly. Poor growth was observed when crude extract from a high rebaudioside-containing stevia (*Stevia rebaudiana* Bertoni) line - an artificial sweetener - was used. Leaf tissue from the control did not show any endopolyploidy but low levels of endopolyploidy (8C) were detected in some treatments.

Keywords: agar, alternative gelling agents, milk, oatmeal agar, table sugar.

INTRODUCTION

Driving down the cost of plant tissue culture is a constant objective of companies and research centers in developing countries (Purohit et al. 2011). Prakash et al. (2004) estimated that chemicals in micropropagation media cost slightly less than 15% of the total cost while the cost of gelling agent per unit media is 73.53%, although that estimate was made in India and is likely to vary from country to country. Alternative low-cost gelling agents have been tested, including sago powder, isabgol husk, guar gum, cassava flour, xantham gum (Maliro and Lameck 2004; Jain and Babbar 2005, 2006; Gour and Kant 2011) and several other media alternatives tested on a hybrid orchid, *Cymbidium* (Teixeira da Silva and Tanaka 2009; Van et al. 2012), papaya and chrysanthemum (Teixeira da Silva 2014a) and also tested on *Syngonium podophyllum* L. (arrowhead vine) in this study. The choice of gelling agent is an important factor determining the success of a plant tissue culture protocol (reviewed by Cameron 2008). The rheological and diffusion properties of gelling agents have been shown to affect shoot development in apple and black locust (Dobrąnszki et al. 2011) and *Amelanchier canadensis* (Fira et al. 2013), and also influence the incidence of hyperhydricity in *Aloe* (Ivanova and van Staden 2011).

Despite the established tissue culture of *S. podophyllum*, either through somatic embryogenesis

(Zhang et al. 2006) or protocorm-like bodies (PLBs; Cui et al. 2008), most commercial tissue culture of arrowhead vine involves the use of shoot tips as clonal propagation units. The use of table sugar or alternative medium additives and gelling agents in the tissue culture of this ornamental and is the fundamental objective of this study. As a practical objective, the ability to identify suitable and effective alternative media additives and gelling agents could potentially allow costs to be reduced.

The objective of this study was thus to test a range of medium additives, some traditional, others not, that might improve shoot and plantlet growth of this ornamental *in vitro*. These additives formed three groups: (i) gelling agents; (ii) liquids serving as alternatives to water; (iii) table sugar versus regular sucrose. There are two reasons to test these substances: (i) as a purely scientific exploration of substances that have never been tested before in the plant science literature; (ii) to find alternatives that could serve as low-cost options for developing countries.

MATERIALS AND METHODS

Chemicals and reagents

All chemicals and reagents, which were of tissue culture grade or the highest grade available, were purchased from Nacalai-Tesque (Kyoto, Japan), Wako Pure Chemical Industries (Osaka, Japan) or Sigma-Aldrich (St.

Louis, MI, USA), whichever offered the best price at the same grade. Gellan gum (Gelrite[®]) was purchased from Merck (USA). The following were bought from local supermarkets in Takamatsu (Japan) (brand name indicated in parentheses): low and full fat milk (Meiji), Coca-Cola[®], instant, granulated coffee (Nescafé, Nestlé), Japanese green tea (Itoh), Oolong tea (Suntory), Darjeeling tea (Tetley), table sugar (Mitsui Sugar Co. Ltd.), corn starch (A-Price). Stevia (*Stevia rebaudiana* Bertoni) was a Chinese cultivar (high level of rebaudioside) grown under field conditions.

The rationale for the selection of these alternatives is explained next. Milk is a daily commodity around the world that is often discarded in a bid to manipulate milk prices (Food and Water Watch 2010). Milk can thus be a suitable substrate if water is not available. Tea and coffee are well-known antioxidants (e.g., Anissi et al. 2014), and thus by removing reactive oxygen species (ROS) (Korir et al. 2014) from in vitro cultures, their growth may be theoretically improved. The exogenous addition of antioxidants has already been shown to improve the growth of hybrid *Cymbidium* (Teixeira da Silva 2013) and *Brassica napus* (Hoseini et al. 2014) cultures. Stevia is an alternative sweetener to sucrose, being as much as 300-fold sweeter (Meireles et al. 2006). To date, no study has yet explored the extract of this plant as an alternative to sucrose in plant in vitro culture. A cheap and popular drink available globally (every country except for Cuba and North Korea) is Coca-Cola[®] (<http://en.wikipedia.org/wiki/Coca-Cola>), which could serve, in a degassed form, as an alternative to sucrose in plant in vitro cultures.

Plant cultures, explants and general media preparation

Original in vitro plant cultures and mother (donor) plants (*S. podophyllum* cv. ‘White Butterfly’) were obtained courtesy of Mr. Kobayashi of the Kobayashi Botanical Gardens (Kan-onji, Kagawa Prefecture, Japan). Plants were established and amassed over several months and maintained as for *Spathiphyllum*, explained in detail in Teixeira da Silva et al. (2006). Initial stock shoots (rootless, 5 cm in length) were transferred to 80 mL of Hyponex[®] (6.5: 6: 19; 3 g/L; Hyponex, Osaka, Japan) medium solidified with 2 g/L of Gellan gum, at 5 shoots/bottle, with 3% (w/v) sucrose in an air-tight Magenta glass bottle (75 mm wide × 250 mm tall) to multiply stocks. Apical shoots 2 cm long with two fully open/developed leaves were harvested from actively growing clonal cultures and used as the explants for all tissue culture experiments. All media were adjusted to pH 5.8 with 1N NaOH or 1 N HCl prior to autoclaving at 121°C, 100 KPa for 15 min. No plant growth regulators were used and additives (gelling agents or medium additives), which were added prior to autoclaving, were not filter sterilized. Shoots from all controls and treatments were cultured at 25 ± 1°C under a 16-h photoperiod with a light intensity of 45 μmol m⁻² s⁻¹ provided by plant growth fluorescent lamps (Plant Lux, Toshiba Co., Japan). All cultures were initiated simultaneously (within 3 days) and tracked for 90 days.

Substrate, medium additive and photoautotrophic treatments

Three sets of experiments were established to test the effect of substrate and/or gelling agent and medium additives on shoot growth and plant development. All experiments were repeated in triplicate. In all experiments, 5 shoots were placed in each 500 mL Magenta bottle and grown on 100 mL of Gellan gum (2 g/L)-solidified Hyponex[®]. In set 1, there was no “standard” control (i.e., treatment against which all other treatments were compared) since no industry standard exists yet; thus treatments were compared against each other. In set 2, distilled water-based medium served as the “standard” or control. In set 3, the “standard” (control) was commercial sucrose.

Set 1 (alternative medium gelling agents). Oatmeal agar (30 g/L), agar, bacto agar, potato dextrose agar and phytigel (all four at 8 g/L), barley starch and corn starch (both at 4 g/L), and Gellan gum (Gelrite[®]) (2 g/L). Pre-experimental trials were conducted to test which concentration was firm, not rock-hard, nor jelly-like so as to support the growth of shoots and plantlets.

Set 2 (innovative additives). Full and low-fat milk (100%, v/v), Oolong tea (50%, v/v), Coca-Cola[®], coffee, stevia extract, Japanese green, Oolong and Darjeeling teas (2%, v/v). Coca-Cola[®] was shaken for 24 h prior to use in medium to de-gas the soft drink. A stock solution of teas, stevia extract and coffee was prepared. For coffee and stevia extract, 10 g/L (coffee powder or fresh stevia leaves) was added to boiling water, stirred well for 5 min and then added to culture medium until a 2% (v/v) final concentration was obtained. Similarly, Japanese green tea and Darjeeling tea stocks were prepared by infusing 5 teaspoons of dry leaves or a tea-bag (approx. 5 g), respectively for 3 min in boiling water and then added to culture medium until 2% (v/v) final concentration was obtained. Oolong tea was purchased as a ready-made bottled tea and was used as such in media.

Set 3 (tissue culture grade sucrose vs table sugar). Table sugar was added to medium at the same concentration as sucrose, i.e., 2% (v/v). Sucrose costs 2200 JPN Yen/kg while refined table sugar (sugar-cane-derived) costs 158 JPN Yen/kg (118 JPN Yen = 1 US\$; January, 2015), i.e., ~14 times higher cost.

Morphological and physiological analyses

The growth and development of plants were evaluated after 90 days following culture initiation, i.e., plating shoot tips (Figure 1A). Survival percentage, total number of newly formed leaves (i.e. excluding the initial two leaves), and total plantlet net fresh weight (FW) were determined. Plantlet survival was calculated based on the percentage of plants being totally green and with root formation. Plantlet height was not measured because of vertical and horizontal expansion of shoot clusters. Chlorophyll content of three random leaves of plantlets from each flask, while still attached to the plant, was measured by a chlorophyll meter (SPAD-502, Minolta Co., Japan) and reported as the SPAD value. The SPAD value is highly correlated with chlorophyll content ($R^2 = 0.89$) and thus serves as a simple and effective way, through SPAD units, to reflect chlorophyll content (Coste et al. 2010).




Figure 1. In vitro micropropagation of *Syngonium podophyllum* L. cv. 'White Butterfly'. Initial explants approximately 2 cm long and with two small, fully-developed leaves (A) were grown on several media, including, 2 g/L Gellan Gum-gelled medium (B), 8 g/L agar-gelled medium (C), 100% (v/v) full fat milk, which curdles and hardens after autoclaving (D), and de-gassed Coca-Cola® on 2 g/L Gellan Gum-gelled medium (E), all supplemented with 3 g/L Hyponex® (6.5: 6: 19) and 3% (w/v) sucrose, demonstrating the capacity to grow (or not) on media with different gelling agents/bases and additives, albeit with different growth responses.

Flow cytometry

The protocol follows that used by Teixeira da Silva and Tanaka (2006), but was adjusted for arrowhead vine. Nuclei were isolated from 0.5 cm² of leaf material (leaf stalk, lamina and mid-vein, pooled) from all treatments in Table 1 by chopping in a few drops (5-10) of nucleic acid extraction buffer (PartecCystain UV Precise P, Germany), and left to digest on ice for 5 min. Three separate leaves (youngest, developing leaves) from three different plants for each treatment were used. The nuclear suspension was then filtered through a 30 µm mesh size nylon filter (CellTrics®) and five times of Partec Buffer A (2 µg/mL 4,6-diamidino-2-phenylindole (DAPI), 2 mM MgCl₂, 10 mM Tris, 50 mM sodium citrate, 1% PVP K-30, 0.1% Triton-X, pH 7.5; Mishiba and Mii 2000) was added at room temperature for 5 min. Nuclear fluorescence was immediately measured using a Partec® Ploidy Analyser and relative fluorescence intensity of the nuclei was analyzed when the coefficient of variation was < 3% by counting a minimum of 5000 nuclei for each sample. Diploid barley (*Hordeum vulgare* L.) cv. 'Ryufu', courtesy of Prof. Shin Taketa (Faculty of Agriculture, Kagawa University), served as the internal control.

Statistical analyses

Experiments were organized according to a randomized complete block design (RCBD, n = 60 per treatment). Each experiment was repeated three times and each experiment had three replicates. For all parameters tested, data analyses were carried out using IRRISTAT version 3.0. Following one-way analysis of variance (ANOVA), Duncan's multiple range test (DMRT) at $P = 0.05$ and the student's t -distribution (standard error, Excel 2010) were used to test for differences between means. Percentage values were arcsin transformed prior to analysis.

RESULTS AND DISCUSSION

Only two studies exist on the tissue culture of *Syngonium podophyllum* (Zhang et al. 2006; Cui et al. 2008). Those studies are important because they indicate protocols for callus formation from somatic embryos and plantlet regeneration from PLBs, respectively, although the terminology pertaining to PLBs in the latter study is being questioned (Teixeira da Silva 2014b). In clonal commercial production, shoot tips are used. This implies that there is a large gap in our understanding of the response of this ornamental in vitro. Thus, by understanding the response of 'White Butterfly', a popular cultivar, to different gelling agents and media additives would narrow this information gap. This study examined the impact of a wide range of substrates and medium additives using the rationale outlined above.

Syngonium shoots grew best on Gellan gum-based medium (Figure 1B), followed by agar-based medium (Figure 1C) but performed poorly on all other agar types. Plantlets performed poorly on medium when supplemented with milk (full-fat or low-fat) (Figure 1D), Coca-Cola® (Figure 1E), coffee or tea (Japanese green, Oolong and Darjeeling teas). The exact reason for this poor performance is not known. However, these additives are not naturally occurring and are man-made, thus one possibility is that enzymes required for their degradation and subsequent uses simply do not exist in plants. For example, in humans, enzymes such as lactase, amylase or catalase (non-exhaustive list) are produced that result in the break-down of lactose, amylose or hydrogen peroxide in milk. However, while catalase is produced by plants, lactase and amylase are not, or are rare (e.g., Stano et al. 2011), thus growth in milk might thus be inhibited simply because the plant cannot degrade the substrate to a

physiologically useful form. One way of being able to use milk by plants would be to genetically modify the plant with a lactase or catalase gene. Observing trends across treatments for gelling agent (set 1) in Table 1, the following can be concluded: (i) plantlet survival and SPAD value were highest when agar or Gellan gum were used; (ii) most number of new leaves and greatest plantlet FW were obtained when Gellan gum served as the gelling agent; (iii) endopolyploidy was detected in cultures with barley or corn starch or potato dextrose agar. Generally, for the clonal production of plants, changes to morphology, genetics or other performance-related indices are undesired. Therefore, for clonal production, the incidence of endopolyploidy may result in undesirable phenotypes. However, in contrast, in ornamentals, where mutations or induced variations are desired to create novel phenotypes, endopolyploidy may serve a useful function. A wider discussion of endopolyploidy may be found in a recent review by Scholes and Paige (2015). Observing trends across treatments for medium additives (set 2) in Table 1, the following can be concluded: (i) all parameters (plantlet survival, plantlet FW, SPAD value, most new leaves formed) were highest when distilled water served as the basal medium liquid; (ii) endopolyploidy, which is correlated to systematics, organ, life strategy and genome size (Barow and Meister 2003), was detected in cultures

grown in the presence of Coca-Cola® or coffee. Finally, in set 3, table sugar could be used as efficiently as tissue culture grade sucrose, resulting in significantly equivalent performance for all parameters observed.




Figure 2. Flow cytometric analysis showing low levels of endopolyploidy (8C) in leaves of *Syngonium podophyllum* L. cv. ‘White Butterfly’ exposed to treatments marked with * in “Leaf ploidy” column of Table 1. 2Cb = 2C peak for diploid barley (*Hordeum vulgare* L.) cv. ‘Ryufu’ (internal standard).

Table 1. Effect of alternative gelling agents, medium additives and carbohydrate sources on *Syngonium podophyllum* L. cv. ‘White Butterfly’ plantlet growth and development, measured 90 days after culture of shoot tips.

Treatments	Plantlet survival (%)	SPAD value	No. new leaves	Plantlet net FW (mg)	Leaf ploidy (2C: 4C: 8C)
<i>Gelling agent</i> ¹					
Agar	91 a	41.6 a	5.2ab	462 b	90 : 10 : 0
Bacto agar	24 cd	36.2 b	1.3 cd	117 d	92 : 8 : 0
Barley starch	31 c	34.6 b	1.1 cd	186 c	90 : 9 : 1*
Corn starch	26 cd	28.0 bc	2.4c	133 d	89 : 8 : 3*
Gellan gum (Gelrite®)	100 a	43.8 a	6.3 a	581 a	88 : 12 : 0
Phytigel	18 d	29.6 bc	4.1 b	126 d	92 : 8 : 0
Potato dextrose agar	0 e	0 d	0.1d	Neg.	93 : 6 : 1*
Oatmeal agar	46 b	24.0 c	0.2d	23 e	91 : 9 : 0
<i>Alternative liquid-based medium additives</i> ¹					
Distilled water-based	100 a	44.2 a	6.4 a	572 a	89 : 11 : 0
Coca-Cola®	0 e	0 d	0 d	Neg.	93 : 6 : 1*
Coffee	0 e	0 d	0 d	Neg.	94 : 5 : 1*
Milk					
Full-fat	62 c	16.1 c	1.4 a	253 d	95 : 5 : 0
Low-fat	58 d	18.2 c	2.1 bc	286 d	95 : 5 : 0
Stevia extract	64 c	31.6 b	3.4 b	351 c	91 : 9 : 0
Tea					
Darjeeling	71 bc	33.2 b	3.2 b	401 b	94 : 6 : 0
Japanese green	77 b	31.6 b	2.4 bc	427 b	93 : 7 : 0
Oolong	70 bc	29.8 b	2.2 bc	416 b	94 : 6 : 0
<i>Carbohydrate source</i> ¹					
TC grade sucrose	100 a	44.4 a	5.6 a	546 a	91 : 9 : 0
Table sugar	98 a	43.6 a	5.1 a	532 a	90 : 10 : 0

Note: ¹ Exact concentrations and methods of preparation are explained in detail in the materials and methods section. Means within a column within each set of experiments (A, B, C) and for each plant character followed by the same letters are not significantly different at $P=0.05$ by Duncan’s multiple range test, $n=60$ per treatment. In all cases, 5 shoot tips were plated per culture vessel. Percentage values arc-sin transformed prior to analysis. FW = fresh weight that exceeds the fresh weight of a single “starting” explant (Figure 1A). Neg. = negligible (< 5 mg). Leaf ploidy levels involved the average of three samples (different leaves) per treatment. * indicates the presence of endopolyploidy (see Figure 2). TC = tissue culture.

Most of the factors assessed in this study have been shown to affect the growth of plantlets in vitro of other species, but never all for a single plant. The choice of gelling agent affected organogenesis in hybrid *Cymbidium* plantlet cultures in which Gellan gum resulted, in general, in better plant growth parameters than Bacto agar and oatmeal agar while the number of roots was highest on Gellan gum as was the fresh and dry mass of shoots and roots although more leaves were produced on Bacto agar (Van et al. 2012). The difference in performance, as indicated above for milk, may be related to the lack of suitable enzymes to degrade the substrates into utilizable carbohydrate sources. Gellan gum is a highly pure gelling agent, thus the existence of other impurities in other gelling agents may also be an influential factor. It is for this reason that Gellan gum tends to be more expensive than regular agar, although prices differ widely between manufacturers. In that study, the chlorophyll content of *Cymbidium* plantlets grown in oatmeal agar was lowest among all basal medium treatments; finally, oatmeal agar-based medium strongly inhibited the initiation of new leaves and roots compared to other gelling agents. As observed in this study (Table 1), Gellan gum formed more PLBs than oat meal agar and potato dextrose agar in another hybrid *Cymbidium* (Teixeira da Silva and Tanaka 2009). Thus, in this study, none of the alternative gelling agents tested could support the growth of arrowhead vine as effectively as Gellan gum. However, in countries where only such gelling agents exist, growth is possible, but may account for poor visual aspect or low productivity. Visual aspect is important for this ornamental, which is prized for its visually attractive leaves, but the visual aspect of poor in vitro growth might not be an important issue if such media were to be used for medicinal plants, horticultural crops or agronomic species.

The choice of gelling agent affected adventitious shoot regeneration capacity and water content (i.e., the state of hyperhydricity) of French marigold (*Tagetes minuta*) shoots (Modi et al. 2009) or *Aloe* (Ivanova and van Staden 2011). The higher the agar concentration, the lower the hyperhydricity of *Dianthus caryophyllus* shoots (Casanova et al. 2008). When phytigel was used as the gelling agent, shoots of *Malus x domestica* (apple) (Turner and Singha 1990), *Pyrus communis* (pear) (Kadoka and Niimi 2003) and *Scrophularia yoshimurae* (Tsay et al. 2006) became hyperhydric, but the physiological status of apple and black locust shoots was attributed to the rheological and diffusion properties of the gelling agents (Dobrąnszki et al. 2011). Van et al. (2012) did not observe hyperhydricity in *Cymbidium* plantlets in any gel-based media. In this study, hyperhydricity was not observed but leaf margins tended to appear "burnt", or dehydrated, possibly due to a lack of moisture in the medium. Seven commercial agar brands caused different organogenic responses in rose (*Rosa hybrid* L. cv. 'Motrea'), lily (*Lilium longiflorum* cv. 'Enchantment') and cactus (*Sulcorebutia alba*) (Scholten and Pierik 1998). *Phalaenopsis* leaf segments obtained from shoots derived from flower-stalk cuttings cultured in vitro on a Gelrite[®]-solidified medium resulted in the formation of more callus-derived PLBs than when agar was used as the medium solidifying agent (Ishii et al. 1998).

Three-fold higher dry weight was observed with tobacco and wild carrot cultures grown on medium gelled with corn starch than on medium gelled with agar (Henderson and Kinnersley 1988). A mixture of corn starch and Gelrite[®] was a suitable substitute for agar in the in vitro cultivation of apple and red raspberry (Zimmerman et al. 1995). Unlike for arrowhead vine (Table 1), starches from barley, corn, potato, rice and wheat were all suitable substitutes for agar in the culture of barley (*Hordeum vulgare* L.) seeds, the most effective being that from barley (Sorvari 1986). 'Isugol', derived from the mucilaginous husk derived from the seeds of *Plantago ovata*, is used in the tissue culture and seed germination of *Syzygium cumini*, *Datura innoxia* (Babbar and Jain 1998), and *Dendrobium chrysotoxum* (Jain and Babbar 2005). The choice of gelling agent affected the regeneration efficiency on selective medium in tulip (*Tulipa* sp.), gladiolus (*Gladiolus* sp.) and tobacco transformation experiments (Chauvin et al. 1999). The level of impurities within a gelling agent might contribute to the outcome of an organogenic pathway, as was demonstrated for *Ranunculus asiaticus* shoots grown in basal medium containing one of three commercial agars (Beruto and Curir 2006). Agar was better than sago powder, guar gum or isabgol husk as a media-solidifying agent for *Balanites aegyptiaca* and *Phyllanthus emblica* (Gour and Kant 2011).

Similarly, where pure water might not be available, or where water may be more expensive than milk, or Coca-Cola[®], or rare, such as in countries with desert conditions or limited water supply, low-fat milk can support the growth of arrowhead vine, but not as effectively as a water-based medium. The prices of water - and expensive but essential commodity - due to its link to so many sectors, will undoubtedly increase (Campbell and Tilley 2014), and this may signal the need to find an alternative to water for plant tissue culture systems in the future.

There is always interest in finding alternatives to sucrose as a carbohydrate source for the heterotrophic culture of plants in vitro. In this study, the use of stevia extract or Coca-Cola[®] as an alternative sugar (carbohydrate) source resulted in poor plantlet growth but table sugar performed as effectively as commercial sucrose (Table 1). In chrysanthemum, other carbohydrates can influence rhizogenesis, caulogenesis (shoots) and somatic embryogenesis (reviewed in Teixeira da Silva et al. 2013), callus and PLB formation in *Cymbidium* hybrids (Teixeira da Silva et al. 2007) and root induction in tree peony (*Paeonia suffruticosa* Andr.) (Wang et al. 2012). There is a possible weakness of this study related to the level at which these liquid additives were added to basal medium and to their osmotic nature. It is possible that tea, at lower concentrations (or higher dilutions), the use of other commercial sodas, or the use of a more dilute concentration of stevia leaf extract may serve as useful forms of carbohydrate and antioxidants in low concentrations for the improved in vitro culture of plants. Evidently, this will require more extensive testing and a wider selection of plants to ascertain the optimal concentrations for practical use in plant tissue culture. If the new substrate has too much of an osmotic gradient, the plant may be stressed as

water flows out from the plant into the medium, thus these substrates and alternatives to plant tissue culture may also serve as new forms of stressors for the study of plant stress physiology. Also, as suggested above, especially for additives like milk, there may be interest in generating transgenic plants carrying the lactase gene to be able to degrade lactose, thus allowing the plant to use this carbohydrate source for growth and development.

ACKNOWLEDGEMENTS

The author thanks Mr. Kobayashi of the Kobayashi Botanical Gardens (Kan-onji, Kagawa Prefecture, Japan) for providing the original *in vitro* stock plants. Sincere thanks to Prof. Michio Tanaka for providing the laboratory facilities to conduct this study. Thanks, too to Dr. Zai-bin Hao (Northern Agricultural University, Harbin, China) for providing fresh stevia plant material. Finally, thanks to Prof. Shin Taketa (Faculty of Agriculture, Kagawa University) for providing barley seeds that served as the internal standard for flow cytometry.

DISCLAIMER AND CONFLICTS OF INTEREST

The author does not specifically endorse any of the brands used in this study. These were used with the pure intention of scientific exploration in mind. The author declares no conflicts of interest, financial or other.

REFERENCES

- Anissi J, El Hassouni M, Ouardaoui A, Sendide K. 2014. A comparative study of the antioxidant scavenging activity of green tea, black tea and coffee extracts: a kinetic approach. *Food Chem* 150: 438-447.
- Babbar SB, Jain N. 1998. 'Subgol' as an alternative gelling agent in plant tissue culture. *Plant Cell Rep* 17: 318-322.
- Barow M, Meister A. 2003. Endopolyploidy in seed plants is differently correlated to systematics, organ, life strategy and genome size. *Plant, Cell Env* 26: 571-584.
- Beruto M, Curir P. 2006. Effects of agar and gel characteristics on micropropagation: *Ranunculus asiaticus*, a case study. In: Teixeira da Silva JA (ed) *Floriculture, Ornamental and Plant Biotechnology: Advances and Topical Issues* (1st ed, Vol. II), Global Science Books Ltd., Isleworth, UK.
- Cameron SI. 2008. Plant tissue culture gelling agents and supports: History, development and function. In: Teixeira da Silva JA (ed) *Floriculture, Ornamental and Plant Biotechnology: Advances and Topical Issues* (1st ed, Vol. V), Global Science Books Ltd., Isleworth, UK.
- Campbell ET, Tilley DR. 2014. The eco-price: how environmental energy equates to currency. *Ecosystem Services* 7: 128-140.
- Casanova E, Moysset L, Trillas MI. 2008. Effects of agar concentration and vessel closure on the organogenesis and hyperhydricity of adventitious carnation shoots. *Biol Plant* 52: 1-8.
- Chauvin J-E, Marhadour S, Cohat J, Le Nard M. 1999. Effects of gelling agents on *in vitro* regeneration and kanamycin efficiency as a selective agent in plant transformation procedures. *Plant Cell Tiss Organ Cult* 58: 213-217.
- Coste S, Baraloto C, Leroy C, Marcon E, Renaud A, Richardson AD, Roggy J-C, Schimann H, Uddling J, Hérault B. 2010. Assessing foliar chlorophyll contents with the SPAD-502 chlorophyll meter: a calibration test with thirteen tree species of tropical rainforest in French Guiana. *Ann. For. Sci.* 67: # 607 (5 pp.)
- Cui J, Liu J-X, Deng M, Chen J-J, Henny RJ. 2008. Plant regeneration through protocorm-like bodies induced from nodal explants of *Syngonium podophyllum* 'White Butterfly'. *HortScience* 43: 2129-2133.
- Dobránszki J, Magyar-Tábori K, Tombácz E. 2011. Comparison of the rheological and diffusion properties of some gelling agents and blends and their effects on shoot multiplication. *Plant Biotechnol Rep* 5: 345-352.
- Fira A, Magyar-Tábori K, Hudák I, Clapa D, Dobránszki J. 2013. Effect of gelling agents on *in vitro* development of *Amelanchier canadensis* 'Rainbow Pillar'. *Int J Hort Sci* 19 (3-4): 76-79.
- Food and Water Watch. 2010. Consolidation and price manipulation in the dairy industry. <http://documents.foodandwaterwatch.org/doc/DairyCompetition-web.pdf> [20 January 2015]
- Gour VS, Kant T. 2011. Efficacy of low cost gelling agents and carbon source alternatives during *in vitro* rooting of *Balanites aegyptiaca*. *Tree For Sci Biotechnol* 5 (1): 58-60.
- Henderson WE, Kinnersley AM. 1988. Corn starch as an alternative gelling agent for plant tissue culture. *Plant Cell Tiss Organ Cult* 15: 17-22.
- Hoseini M, Ghadimzadeh M, Ahmadi B, Teixeira da Silva JA. 2014. Effects of ascorbic acid, alpha-tocopherol and glutathione on microspore embryogenesis in *Brassica napus* L. *In Vitro Cell Dev Biol - Plant* 50 (1): 26-35.
- Ishii Y, Takamura T, Goi M, Tanaka M. 1998. Callus induction and somatic embryogenesis of *Phalaenopsis*. *Plant Cell Rep* 17: 446-450.
- Ivanova M, van Staden J. 2011. Influence of gelling agent and cytokinins on the control of hyperhydricity in *Aloe polyphylla*. *Plant Cell Tissue Organ Cult* 104: 13-21.
- Jain R, Babbar SB. 2005. Guar gum and isubgol as cost-effective alternative gelling agents for *in vitro* multiplication of an orchid, *Dendrobium chrysotoxum*. *Curr Sci* 88: 292-295.
- Jain R, Babbar SB. 2006. Xanthan gum: An economical substitute of agar for *in vitro* multiplication of an orchid, *Dendrobium chrysotoxum* Lindl. *Curr Sci* 91: 27-28.
- Kadoka M, Niimi Y. 2003. Effects of cytokinin types and their concentration on shoot proliferation and hyperhydricity in *in vitro* pear cultivar shoots. *Plant Cell Tiss Organ Cult* 72: 261-265.
- Korir MW, Wachira FN, Wanyoko JK, Ngure RM, Khalid R. 2014. The fortification of tea with sweeteners and milk and its effect on *in vitro* antioxidant potential of tea product and glutathione levels in an animal model. *Food Chem* 145: 145-153.
- Maliro FAM, Lameck G. 2004. Potential of cassava flour as a gelling agent in media for plant tissue cultures. *African J Biotechnol* 3: 244-247.
- Meireles MAA, Wang G-M, Hao Z-B, Shima K, Teixeira da Silva JA. 2006. Stevia (*Stevia rebaudiana* Bertoni): futuristic view of the sweeter side of life. In: Teixeira da Silva JA (ed). *Floriculture, Ornamental and Plant Biotechnology: Advances and Topical Issues* (1st ed., Vol. IV), Global Science Books, Ltd., Isleworth, UK, pp. 416-425.
- Mishiba K, Mii K. 2000. Polysomaty analysis in diploid and tetraploid *Portulaca grandiflora*. *Plant Sci* 156: 213-219.
- Modi P, Sinha A, Kothari SL. 2009. Reduction of hyperhydricity in micropropagated French marigold (*Tagetes patula* L.) plants by modified medium parameters. *Floriculture Ornamental Biotechnol* 3: 40-45.
- Prakash S, Hoque MI, Brinks T. 2004. Culture media and containers. In: Low cost options for tissue culture technology in developing countries, FAO/IAEA Division of Nuclear Techniques in Food and Agriculture, Vienna, pp. 29-40.
- Purohit SD, Teixeira da Silva JA, Habibi N. 2011. Current approaches for cheaper and better micropropagation technologies. *Int J Plant Dev Biol* 5: 1-36.
- Scholes DR, Paige KN. 2015. Plasticity in ploidy: a generalized response to stress. *Trends Plant Sci* [in press]. doi: 10.1016/j.tplants.2014.11.007
- Scholten HJ, Pierik RLM. 1998. Agar as a gelling agent: Chemical and physical analysis. *Plant Cell Rep* 17: 230-235.
- Sorvari S. 1986. The effect of starch gelatinized nutrient media in barley anther cultures. *Ann Acad Sci Fenn* 25: 127-133.
- Stano J, Siekel P, Neubert K, Mi ieta K. 2011. Study of gherkin lactase in cell culture and in seedlings. *Acta Histochem* 113 (6): 631-635.
- Teixeira da Silva JA, Giang DTT, Chan M-T, Sanjaya, Norikane A, Chai M-L, Chico-Ruiz J, Penna S, Granström T, Tanaka M. 2007. The influence of different carbon sources, photohetero-, photoauto- and

- photomixotrophic conditions on protocorm-like body organogenesis and callus formation in thin cell layer culture of hybrid *Cymbidium* (Orchidaceae). *Orchid Sci Biotechnol* 1 (1): 15-23.
- Teixeira da Silva JA, Giang DTT, Tanaka M. 2006. Novel photoautotrophic micropropagation of *Spathiphyllum*. *Photosynthetica* 44 (1): 53-61.
- Teixeira da Silva JA, Shinoyama H, Aida R, Matsushita Y, Raj SK, Chen F. 2013. Chrysanthemum biotechnology: *Quo vadis?* *Critical Rev Plant Sci* 32 (1): 21-52.
- Teixeira da Silva JA, Tanaka M. 2006. Multiple regeneration pathways via thin cell layers in hybrid *Cymbidium* (Orchidaceae). *J Plant Growth Reg* 25 (3): 203-210.
- Teixeira da Silva JA, Tanaka M. 2009. Impact of gelling agent and alternative medium additives on hybrid *Cymbidium* protocorm-like body and callus formation. *Floriculture Ornamental Biotechnol* 3: 56-58.
- Teixeira da Silva JA. 2013. Impact of paper bridges, activated charcoal, and antioxidants on growth and development of protocorm-like bodies of hybrid *Cymbidium*. *In Vitro Cell Dev Biol Plant* 49 (4): 414-420.
- Teixeira da Silva JA. 2014a. Novel factors affecting in vitro shoot culture of chrysanthemum (*Dendranthema x grandiflora* (Ramat.) Kitamura). *Bot Lithuanica* 20 (1): 27-40.
- Teixeira da Silva JA. 2014b. Should the term protocorm-like body be used exclusively for orchids? *J Plant Dev* 21: 161-166.
- Tsay H-S, Lee C-Y, Agrawal DC, Basker S. 2006. Influence of ventilation closure, gelling agent and explant type on shoot bud proliferation and hyperhydricity in *Scrophularia yoshimurae* - a medicinal plant. *In Vitro Cell Dev Biol - Plant* 42: 445-449.
- Turner SR, Singha S. 1990. Vitrification of crab apple, pear and gum on gellan gum-solidified culture medium. *Hort Sci* 25: 1648-1650.
- Van PT, Teixeira da Silva JA, Tanaka M. 2012. How does choice of substrate and culture conditions affect the growth and development of *Cymbidium* cv. Green Planet 'Energy Star' protocorm-like bodies? *Eur J Hort Sci* 77 (5): 219-225.
- Wang H-Y, He S-L, Tanaka M, Pham TV, Teixeira da Silva JA. 2012. Effect of IBA concentration, carbon source, substrate, and light source on root induction ability of tree peony (*Paeonia suffruticosa* Andr.) plantlets in vitro. *Eur J Hort Sci* 77 (3): 122-128.
- Zhang Q, Chen J-J, Henny RJ. 2006. Regeneration of *Syngonium podophyllum* 'Variegatum' through direct somatic embryogenesis. *Plant Cell Tiss Organ Cult* 84: 181-188.
- Zimmerman RH, Bhardwaj SV, Fordham IM. 1995. Use of starch-gelled medium for tissue culture of some fruit crops. *Plant Cell Tiss Organ Cult* 43: 207-213.