

Short communication: A laboratory study to validate the impact of the addition of *Alnus nepalensis* leaf litter on carbon and nutrients mineralization in soil

GAURAV MISHRA , KRISHNA GIRI, ANTARA DUTTA, SATABDI HAZARIKA, PRATUL BORGHAIN
Rain Forest Research Institute, P.O. Box 136, A.T. Road (East), Jorhat-785001, Assam, India. Tel.: +91-376-2305101, Fax.:+91-376-2305130,
email: gaurav.mishra215@gmail.com

Manuscript received: 15 December 2015. Revision accepted: 31 December 2015.

Abstract. Mishra G, Giri K, Dutta A, Hazarika S, Borgohain P. 2016. A laboratory study to validate the impact of the addition of *Alnus nepalensis* leaf litter on carbon and nutrients mineralization in soil. *Nusantara Bioscience* 8: 5-7. Plant litter or residues can be used as soil amendment to maintain the carbon stock and soil fertility. The amount and rate of mineralization depends on biochemical composition of plant litter. *Alnus nepalensis* (Alder) is known for its symbiotic nitrogen fixation and capability to restore fertility of degraded lands. A laboratory incubation experiment was conducted for 60 days under controlled conditions to validate the carbon and nutrients mineralization potential of alder litter. Soil fertility indicators, i.e. soil organic carbon (SOC), available nitrogen (N), available phosphorus (P), and available potassium (K) were analyzed using standard procedures. Significant differences were observed in the soil properties after addition of litter. Nutrient composition of alder litter was found superior by providing significantly higher organic matter and helped in better nutrient cycling. Therefore, alder based land use system may be replicated in other degraded lands or areas for productivity enhancement which is important for sustaining biodiversity and soil fertility.

Key words: *Alnus nepalensis*, carbon, nutrients, Nagaland

INTRODUCTION

Shifting or Jhum cultivation contributes significantly in alterations of soil properties and diminishes soil quality (Ayoubi et al. 2011). Despite being the site specific nature of this land use pattern, it may also lead to desertification, soil erosion, biodiversity loss, climate change, etc. (Chase and Singh 2014). Concerning Northeast India, Jhum cultivation is predominant in all the seven states that pose a serious threat. This region witnessed drastic land use changes in the recent past which leads to decline in soil fertility and productivity. According to Oliveira et al. (2014), available organic materials can be used as soil amendments to manage the fertility status. Moreover, in context to small hold agriculture of this region, plant residues in the form of litter are the potential nutrient sources. Hence, to cope up with the problem of declining soil quality, people of this region developed some indigenous knowledge based agroforestry systems with the aim of sustaining the system (Ramakrishnan 1992). According to the report of NEPED and IIRR (1999), alder based agroforestry system provides at least 57 food crops to supplement the rice cultivation. *Alnus nepalensis* D. Don, or alder is a non-leguminous nitrogen fixing tree species occurs naturally throughout the Himalayas.

The importance of alder tree was recognized by the tribal communities long back 200 years ago and utilized by the farmers of Angami, Chakhasang, Chang, Yimchaunger and Konyak tribes (Gokhle et al. 1985). Alder is known for its symbiotic relationship with *Frankia* (nitrogen-fixing

actinomycetes) and therefore able to restore fertility status of degraded Jhum lands (Rathore et al. 2010). Moreover, the litter of alder also adds P, K, Ca and other nutrient through the biomass decomposition. Incorporation of alder plant litter in soils can maintain organic carbon (OC) content, improve soil physical properties, enhance biological activities and increase nutrient availability. Therefore, this study was aimed to validate the effect of incorporation of alder leaves on soil organic carbon (SOC) and nutrient content of soil incubated under laboratory conditions.

MATERIALS AND METHODS

The soil used in this study was collected from an abandoned Jhum land of Khonoma District of Nagaland, India. The study site is located at Longitude E95°00'281" Latitude N25°37'688 and an elevation of 1531m above mean sea level. The site was sandy loam in texture. Soil samples were collected from a depth of 30cm from three points. Physicochemical properties of the soil were determined in Soil Testing Laboratory of Rain Forest Research Institute, Jorhat, Assam and initial soil analysis is presented in Table 1. Fresh alder leaves were collected from 10 year old plantation, washed in distilled water, dried at 65°C for 48 h, milled and passed through a 1mm sieve. The incubation study was done as per the methodology given by Abbasi et al. (2011). 100 g soil sample was kept in glass jars and moisture content was maintained at field capacity via addition of distilled water. A total of 15 jars were used (5

Table 1. Physicochemical properties of the soil (0-30 cm) used in the study.

Soil properties	Values
Bulk density	0.80
Silt (%)	26.8
Sand (%)	25.9
Clay (%)	47.2
pH	4.51
CEC (c mol kg ⁻¹)	5.06
OC (%)	3.18
Available N (%)	0.001
Available P (ppm)	9.0
Available K (ppm)	223.00

treatments and 3 replications). The treatments were mixing of dry milled alder leaves in different doses, ie: (i) control; (ii) 2.5 g; (iii) 5 g; (iv) 7.5 g and (v) 10 g. This soil sampling and analysis was repeated after every 15 days. SOC, N, P and K were estimated using previously described standard methodology (Walkley and Black 1934; Subbiah and Asija 1956; Bray and Kurtz 1945; Morwin and Peach 1951). Significance of various parameters were tested by F-tests ($P < 0.05$) and Duncan Multiple Range Test (DMRT) using SPSS version 16.0.

RESULTS AND DISCUSSION

Soil organic matter (SOM) is the main component of soil as it provides major portion of soil carbon pool, supports microbiological diversity and ensures the long term supply of nutrients through decomposition (Woomer et al. 1994). The value of SOC decreased in control as compared other treatments during the study period. Addition of alder leaf litter produced significant effect on SOC content in incubated soil at 30, 45 and 60 days (Figure 1). Among the treatments, after 60 days of incubation the value ranged from a minimum of 2.39% in control and a maximum of 3.96% (Treatment 5). The differences were statistically highly significant at 60 days ($F=23.288$, $p=0.00$), where higher dose of alder leaves was found to be superior in maintaining SOC status in long term basis. In general, the added plant residues increased organic matter stock in soil and thereby increased N mineralization and N transformation processes in soil (Abbasi et al. 2015).

Nitrogen is an important macronutrient and plays major role in plant growth and development, but its availability in soil is a major factor in governing the productivity of any ecosystem (Gairola et al. 2012). Soil N content was found to be more in treatment 5 in all sampling dates. After 15 days of incubation is no significant effect of leaf addition, but after 30 days, N content in all the treatments was significantly superior to control (Figure 2). Sudden increase in N content can be supported by the fact that initial low N availability in soil increased the mineralization of litter. The effect of leaf addition also produces significant effect at 45 and 60 days. After 60 days of addition, effects were




Figure 1. Soil organic carbon (%) under different treatments




Figure 2. Available N (%) under different treatments




Figure 3. Available P (ppm) under different treatments




Figure 4. Available K (ppm) under different treatments

pronounced and value ranged between 0.016% (control) to 0.096% (Treatment 5). Alder leaves provides significant amount of litter and organic matter, as evident by a significant increase in soil N after 60 days ($F=163.152$, $p=0.00$). Our results were in accordance with findings of Tripathi et al. (2009), who reported about the role of alder litter in increasing the nitrogen content of soil. N mineralization is slowed down after 45 days as high N contents in soils slowed down the decomposition rates of litter (Liu et al. 2010).

Like N, P is also equally important for plant growth and development. Huge amount of P is stored in soil in different forms but the major concern regarding the availability of P is fixation (Chatterjee et al. 2014). Among the treatments, no significant differences were observed after 15 days, while the effects are more prominent after 30, 45 and 60 days, respectively (Figure 3). After 60 days of incubation, amount of P ranged from 9.17 ppm (Control) to 10.93 ppm (Treatment 5) and alder leaves contributed significantly in maintaining the P status in comparison to control ($F=6.298$, $p=0.008$). Similarly, K in plants plays a key role in biochemical processes like enzyme activation, synthesis of macromolecules, regulation of transpiration etc (Brady 1996). No significant difference was observed in K content after 15 and 30 days can be supported by the fact that amount of K in soil is directly related to the parent rock material (Figure 4). Moreover, the effects are more and less similar after 45 ($F=3.755$, $p=0.041$) and 60 ($F=5.936$, $p=0.010$) days. After 60 days of incubation, amount of K ranged from 206.00 ppm (Control) to 262.00 ppm (Treatment 5). This is due to the fact that organic matter through its slow decomposition maintains the availability of K (Basumatary and Bordoloi 1992).

In conclusion, the experiment showed that soil amended with alder leaves displayed wide variation of SOC and nutrient mineralization depending on the amount of plant litter added. This study suggested that alder litter showing rapid N mineralization so can be used for early N demands of a crop in degraded lands. Uses of such locally available plant materials in the regions prone to low fertility status may be a useful management strategy for nutrient restoration. Our study provides the scientific validation of alder based system in enhancing the fertility status of abandoned Jhum land. In view of this, there is an urgent need to replicate alder based system for productivity enhancement in Jhum lands. In addition, to mitigate the impacts of climate change, the fallow Jhum lands can be considered as potential areas for alder reforestation for increasing the carbon stock.

ACKNOWLEDGEMENTS

Financial support for this work by the Indian Council of Forestry Research and Education, Dehradun, India, is gratefully acknowledged. The authors declare that there is

no conflict of interests regarding the publication of this paper.

REFERENCES

- Abbasi KM, Tahir MM, Sabir N, Khurshid M. 2015 Impact of the addition of different plant residues on nitrogen mineralization-immobilization turnover and carbon content of a soil incubated under laboratory conditions. *Solid Earth* 6: 197-205.
- Abbasi MK, Hina M, Tahir MM. 2011. Effect of *Azadirachta indica* (neem), sodium thiosulphate and calcium chloride on changes in nitrogen transformations and inhibition of nitrification in soil incubated under laboratory conditions. *Chemosphere* 82: 1629-1635.
- Ayoubi S, Khormali F, Sahrawat KL, Rodrigues de Lima AC. 2011. Assessing impacts of land use change on soil quality indicators in a loessial soil in Golestan Province, Iran. *J Agr Sci Tech* 13: 727-742.
- Basumatary A, Bordoloi PK. 1992. Forms of potassium in some soils of Assam in relation to soil properties. *J Indian Soc Soil Sci* 40(3): 443-446.
- Brady NC. 1996. *The Nature and Properties of Soil*, 10th ed. Prentice Hall, New Delhi
- Bray RH, Kurtz LT. 1945. Determination of total, organic and available forms of phosphorus in soils. *Soil Sci* 59: 39-45.
- Chase P, Singh OP. 2014. Soil nutrients and fertility in three traditional land use systems of Khonoma, Nagaland, India. *Res Environ* 4(4): 181-189.
- Chatterjee D, Datta SC, Manjiaiah KM. 2014. Fractions, uptake and fixation capacity of phosphorus and potassium in three contrasting soil orders. *J Soil Sci Pi Nutr* 14 (3): 640-656.
- Gairola S, Sharma CM, Ghildiyal SK, Suyal S. 2012. Chemical properties of soils in relation to forest composition in moist temperate valley slopes of Garhwal Himalaya, India. *The Environment* 32 (4): 512-523.
- Gokhle AM, Zeliang DK, Kevichusa R, Angami T, Bendangnugsang S. 1985. *The Use of Alder Tree*. State Council of Education Research & Training, Government of Nagaland, Kohima, Nagaland.
- Liu P, Huang J, Sun OJ, Han X. 2010. Litter decomposition and nutrient release as affected by soil nitrogen availability and litter quality in a semiarid grassland ecosystem. *Oecologia* 162: 771-780.
- Morwin HD, Peach PM. 1951. Exchangeability of soil potassium in and, silt and clay fractions as influenced by the nature of complementary exchangeable cations. *Proc Natl Acad Sci USA* 15: 125-128.
- NEPED and IIRR. 1999. *Building Upon Traditional Agriculture in Nagaland, India*. Nagaland Environmental Protection and Economic Development, India and International Institute of Rural Reconstruction, Philippines.
- Oliveira SP, Lacerda NB, Blum SC, Escobar MEO, Oliveira TS. 2014. Organic carbon and nitrogen stocks in soils of northeastern Brazil converted to irrigated agriculture. *Land Degrad Dev* 26: 9-21.
- Ramakrishnan PS. 1992. *Shifting Agriculture and Sustainable Development: An Interdisciplinary Study from Northeastern India*. MAB Book Ser., UNESCO, Paris & Parthenon Publishing Group, Carnforth, Lancs., U.K.
- Rathore SS, Karunakaran K, Prakash B. 2010. Alder based farming system, a traditional farming system in Nagaland for amelioration of Jhum land. *India J Trad Knowl* 9 (4): 677-680
- Subbiah BV, Asija CL. 1956. A rapid procedure for the estimation of available nitrogen in soils. *Curr Sci* 25: 256-260.
- Tripathi OP, Pandey HN, Tripathi RS. 2009. Litter production, decomposition and physico-chemical properties of soil in 3 developed agroforestry systems of Meghalaya, Northeast India. *Afr J Pl Sci* 3 (8): 160-167.
- Walkley A, Black IA. 1934. An examination of the Degtjareff method for determining soil organic matter, and a proposed modification of the chromic acid titration method. *Soil Sci* 37: 29-38.
- Woomer PL, Martin A, Albrecht A, Reseck DVS, Scharpenseel HW. 1994. *The importance and management of soil organic matter in the tropics*. In: Woomer PL, Swift MJ (eds) *The Biological Management of Tropical Soil Fertility*. Wiley, Chichester, UK.