

Short Communication: In vitro response of papaya (*Carica papaya*) to plant growth regulators

JAIME A. TEIXEIRA DA SILVA

Retired/independent scientist, Miki cho post office, Ikenobe 3011-2, P.O. Box 7, Kagawa-Ken, 761-0799, Japan. email: jaimetex@yahoo.com

Manuscript received: 11 April 2016. Revision accepted: 2 May 2016.

Abstract. Teixeira da Silva JA. 2016. *In vitro* response of papaya (*Carica papaya*) to multiple plant growth regulators. *Nusantara Bioscience* 8: 77-82. The use of plant growth regulators (PGRs) in papaya (*Carica papaya* L.) tissue culture is essential for tissue and organ culture in vitro. In this study, in a bid to expand the information available on the response to PGRs, a wide range of PGRs, roughly divided into four groups (auxins, cytokinins, alternative PGRs, growth inhibitors and retardants) was tested. Among them, the auxins 2,4-D, dicamba and picloram formed most callus (hard and soft). Callus inductions by chitosan and coconut water are novel results for papaya. Shoots only formed in response to BA and TDZ, but TDZ-induced shoots were fasciated and/or hyperhydric. These results provide novel perspectives for papaya researchers who may have recalcitrant genotypes or tissues that are unresponsive in vitro.

Keywords: 2,4-D, dicamba, PGR, picloram, thidiazuron

Abbreviations: 2,4-D, 2,4-dichlorophenoxyacetic acid; 2,4,5-T, 2,4,5-trichlorophenoxyacetic acid; 2iP, N^6 -[2-isopentenyl] adenine (syn. 6-aminocaproic acid); ABA, (\pm)-*cis,trans*-abscisic acid; AC, activated charcoal; Ads, adenine hemisulphate; BA, 6-benzyladenine (syn. BAP, 6-benzylaminopurine); BNOA, β -naphthoxyacetic acid; BSSAA, benzoseniyl-3-acetic acid; 4-CPPU, *N*-(chloro-4-pyridyl)-*N*-phenylurea (or forchlorfenuron); dicamba, 3,6-dichloro-2-methoxybenzoic acid (syn. 3,6-dichloro-*o*-anisic acid); GA₃, gibberellic acid; JA, (\pm)-jasmonic acid; Kin, kinetin; MeJa, methyl jasmonate; PGR, plant growth regulator; picloram, 4-amino-3,5,6-trichloro-2-pyridinecarboxylic acid; PPF, photosynthetic photon flux density; SA, salicylic acid; TCL, thin cell layer; TDZ, *N*-phenyl-*N'*-1,2,3-thiadiazol-5-yl-urea or thidiazuron; TIBA, 2,3,5-triiodobenzoic acid; ZR, zeatin riboside (syn. 9-(β -D-ribofuranosyl)-*trans*-zeatin or N^6 -(*trans*-4-hydroxy-3-methyl-2-buten-1-yl)adenosine)

INTRODUCTION

Papaya (*Carica papaya* L.; Caricaceae) is a popular tropical fruit, but also contains important secondary metabolites (Canini et al. 2007). This plant is usually propagated by seed (Teixeira da Silva et al. 2007; Jiménez et al. 2014). Somatic embryos are a stable form of clonal propagation and are useful for the establishment of suspension cultures (Anandan et al. 2012). Consequently, somatic embryogenesis has been well explored in papaya (Teixeira da Silva et al. 2007) and to achieve this, several plant growth regulators (PGRs) have been employed throughout the literature. This will be elaborated upon in more detail in the discussion. A robust regeneration system allows for a greater chance of recovering transgenic plants in vitro (Tennant 2010). By adjusting the ratios of red : blue light-emitting diodes (LEDs), papaya plant growth can be manipulated, and photoautotrophic micropropagation is an excellent means to increase plantlet biomass or to alter plant architecture (Teixeira da Silva 2014a). Teixeira da Silva (2013) noted that cotyledonary tissue exposed to five auxins (2,4,5-trichlorophenoxyacetic acid, 2,4,5-T; indole-3-acetic acid, IAA; indole-3-butyric acid, IBA; NAA; β -naphthoxyacetic acid, BNOA) could form axillary roots. A previous study (Teixeira da Silva 2014b) compared the response of six previously established protocols. In that study, friable callus or embryogenic callus was induced

from cotyledons that were then encapsulated as synthetic seed (synseed). These synseed were then subjected to low-temperature storage and cryopreservation, which are two important techniques for preservation of important horticultural germplasm (Cha-um and Kirdmanee 2007; Engelmann 2011; Sharma et al. 2013).

This study exposed two explants (leaves and stem thin cell layers, or TCLs) of a papaya cultivar to a wide range of PGRs, many of which have not yet been tested in papaya tissue culture, to assess developmental responses. TCLs were selected as they are sensitive tissues useful for plant tissue culture (Teixeira da Silva and Dobránszki 2013a, 2015; Teixeira da Silva et al. 2015).

MATERIALS AND METHODS

All chemicals and reagents were purchased from either Sigma-Aldrich (St. Louis, USA), Wako Chemical Co. (Osaka, Japan), or Nacalai Tesque (Kyoto, Japan).

Seeds of a hybrid papaya (*Carica papaya* L. cv. 'Sunrise Solo') cultivar were surface sterilized and germinated using the protocol of Giang et al. (2011) and Teixeira da Silva (2013, 2014a, 2014b). In brief, seeds were soaked for 48 h then washed in running tap water with scrubbing to remove the sarcotesta. Seeds that passed a floatation test (viable seeds) were surface sterilized in

0.1% HgCl₂ + 2-3 drops of Tween-20 for 5 min, rinsed 3 times in sterilized distilled water (SDW), sprayed with 80% ethanol for 1 min then rinsed 3 times in SDW. Surface-sterilized seeds were slightly embedded (5/Petri dish; As-One, Osaka, Japan) in autoclaved (100 KPa; 21 min) full-strength (macro- and micronutrients) PGR-free Murashige and Skoog (1962) (MS) basal medium at pH 5.8 (adjusted with 1N NaOH or HCl) containing 3% sucrose and 2 g/L gellan gum (Gelrite[®]; Merck, USA). Petri dishes were sealed with Parafilm[®] (Pechiney Plastic Packaging Co., Chicago, USA) and placed at 25°C in the dark for 7 days. Seedlings then transferred to 250-ml Erlenmeyer flasks (5 plantlets per flask) containing 50 ml of Hyponex[®] (N:P:K = 6.5 : 6 : 19; Hyponex Japan Corp., Tokyo, Japan), 3% sucrose and 2 g/L Gelrite[®] and placed under a 16-h photoperiod with a photosynthetic photon flux density (PPFD) of 45 μmol m⁻² s⁻¹ provided by plant growth fluorescent lamps (Plant Lux, Toshiba Co., Japan). Young leaves of one-month-old rooted seedlings established in vitro were used for the experiments that follow. Transverse thin cell layers (tTCLs), which are useful explants for better control of plant organogenesis in vitro (Teixeira da Silva and Dobránszki 2013a, 2014), were prepared from seedling stems, 1 mm thick and 1-2 mm in diameter.

The base of young leaves of two-month-old established in vitro rooted seedlings, including mid-vein tissue, as well as stem tTCLs, were exposed to four groups of PGRs in the light (16-h photoperiod; PPFD = 45 μmol m⁻² s⁻¹) and in the dark. The rationale was to assess whether a dose of any single PGR could induce organogenesis. Four categories of PGRs were tested. Group 1, auxins (3,6-dichloro-2-methoxybenzoic acid (syn. 3,6-dichloro-*o*-anisic acid)), dicamba (pesticide that mimics an auxin); 2,4-D (pesticide that mimics an auxin); 4-amino-3,5,6-trichloro-2-pyridinecarboxylic acid, picloram (a herbicide)). Group 2, cytokinins (adenine hemisulphate, AdS; 6-benzyladenine, BA (syn. 6-benzylaminopurine; see notes in Teixeira da Silva 2012); benzoselenienyl-3-acetic acid, BSSAA; *N*-(chloro-4-pyridyl)-*N*-phenylurea (or forchlorfenuron), 4-CPPU; kinetin, Kin; *N*-phenyl-*N'*-1,2,3-thiadiazol-5-yl-urea or thiazuron, TDZ (cytokinin-like); zeatin riboside (syn. 9-(*-D*-ribofuranosyl)-*trans*-zeatin or *N*⁶-(*trans*-4-hydroxy-3-methyl-2-buten-1-yl)adenosine), ZR; *N*⁶-[2-isopentenyl] adenine (syn. 6-(*2*, -dimethylallylamino) purine), 2iP). Group 3, other PGRs (chitosan (a fungal elicitor); coconut water (CW); gibberellic acid, GA₃; jasmonic acid, JA; methyl jasmonate, MeJA; salicylic acid (SA); 1-triacontanol (TRIA; syn. melissyl alcohol, myricyl alcohol). Group 4, growth inhibitors and retardants ((±)-*cis,trans*-abscisic acid, ABA; ancymidol; chlormequat; mepiquat; paclobutrazol; uniconazole; 2,3,5-triiodobenzoic acid, TIBA). Four concentrations (1, 2, 4 and 8 mg/L) of each PGR was tested separately, except for CW, which was tested at 1, 2, 4 and 8% (v/v). Coconuts were purchased fresh in a green state from Osaka fruit market, transported to Kagawa under cooled conditions. A hole was drilled using aseptically treated tools and CW was poured into a sterile plastic bottle and frozen until needed, or used fresh immediately after extraction. The control was PGR-free MS medium.

The formation of shoots or callus was assessed with the naked eye and by light microscopy and quantified after 60 days. The amount of hard callus (HC), soft, friable callus (SFC) and shoots that formed was not quantified. More discussion about HC and SFC may be found in Teixeira da Silva (2104b).

Experiments were organized according to a randomized complete block design with three blocks of 10 replicates per treatment. All initial trials were repeated once. Only those treatments that showed an organogenic response (Table 1) were repeated three times.

RESULTS AND DISCUSSION

A wide range of PGRs and growth retardants on the organogenesis of papaya in vitro was tested since several of these PGRs have not yet been used in papaya tissue culture. The most popular PGR used in papaya somatic embryogenesis, for example, is 2,4-D, given its success across a wide range of cultivars. The in vitro responses of papaya explants cultured on MS media supplemented with different PGR are given in Table 1. No growth or explant response was observed for any of the group 4 PGRs (growth retardants). The most responsive group was the auxins in group 1, particularly 2,4-D, picloram and dicamba. Other auxins had already been tested in Teixeira da Silva (2013). Both leaf explants and tTCLs were responsive to 2,4-D in the light and in the dark, forming both HC (Figure 1E, F, G, H) and SFC (Figure 1D). HC formed in response to dicamba and picloram, both in the dark, but only in tTCLs in the latter and in leaves and tTCLs in the former while SFC showed a similar trend with respect to explant receptivity as dicamba, but in the light. In group 2, Kin and TDZ formed SFC, but TDZ showed a wider response for both explants and both in the dark and light while 2iP showed a limited formation of HC by leaves in the dark. In group 3, chitosan and GA₃ formed HC, independent of the dark or light conditions, but only tTCLs were responsive for chitosan. CW formed some SFC from tTCLs in both the light and dark. TRIA showed limited HC formation from tTCLs in the dark. Shoots only formed in the presence of BA or TDZ, but those from TDZ tended to be fasciated and/or hyperhydric (Figure 1C). As expected, none of the growth retardants (group 4) induced any response (callus or shoots).

Even though somatic embryogenesis was widely studied in papaya (see below), no clear somatic embryos or somatic embryo-like structures were observed in this study, even within the wide range of PGRs tested. One possible reason may be that this study only tested individual PGRs or a single medium step, while most other studies employed PGR combinations or transfer to more than one medium. Therefore, it is possible that the HC or SFC observed in this study could develop somatic embryos had they been transferred to alternative PGR-containing media for somatic embryo induction. Another reason may be the range of concentrations tested (1-8 mg/L). For example, de Almeida et al. (2001) required a higher concentration of 2,4-D (10 mg/L) to induce friable callus and then, after

Table 1. Effect of plant growth regulators on organogenesis (HC, SFC or shoots) from two-month-old leaves or stem tTCLs of papaya (*Carica papaya* L.) ‘Sunrise Solo’ in the light and dark, assessed after 60 days in culture. All plant growth regulator (PGRs) and concentrations not displayed did not show any growth or organogenic response (even if the fresh weight may have increased).

PGR	HC				SFC				Shoots			
	Light		Dark		Light		Dark		Light		Dark	
	Leaf	Stem tTCL	Leaf	Stem tTCL	Leaf	Stem tTCL	Leaf	Stem tTCL	Leaf	Stem tTCL	Leaf	Stem tTCL
Control No PGRs												
Group 1 2,4-D	1-8	1-8	1-8	1-8	1-8	1-8	1-8	1-8				
Dicamba			1, 2	1, 2	1	1						
Picloram				1		1						
Group 2 AdS												
BA									1, 2	1, 2, 4		
BSSAA												
4-CPPU												
Kin						1						
TDZ					2, 4	1-8	2	1-8			1-8	1-8
ZR												
2iP	1		1									
Group 3 Chitosan		1, 2		1, 2								
CW					1		1					
GA ₃	1	1	1, 2	1, 2								
JA												
MeJA												
SA												
TRIA				1								
Group 4 ABA												
Ancymidol												
Chlormequat												
Mepiquat												
Pacllobutrazol												
Uniconazole												
TIBA												

Callus types: HC, hard callus; SFC, soft, friable callus. Shading key: no shading = no response or only explant expansion; yellow = SFC; orange = HC; green = shoots. Numbers in table boxes: no numbers = no response; all other numbers correspond to either 1, 2, 4 or 8 mg/L of the PGRs, or 1, 2, 4 or 8% of CW. See text for PGR abbreviations

transfer to another medium, somatic embryos.

Apparently lower levels of organogenesis observed in tTCLs relative to conventional leaf explants does not necessarily reflect a poorer performance of tTCLs since organogenesis is restricted by the actual explant size, surface area and volume (Teixeira da Silva 2013; Teixeira da Silva and Dobránszki 2013b). To correct for this difference between the organogenic outcome of a TCL and a conventional explant, the Plant Growth Correction Factor (Teixeira da Silva and Dobránszki 2011, 2014) takes three aspects (explant size, surface area, and volume) into consideration, allowing the organogenic outcome to be directly compared. Another possibility may relate to the timing of sampling and the developmental state of the tissues, both factors playing a considerable outcome on the organogenic response (Teixeira da Silva and Dobránszki 2013b). It is thus a possibility that the observed results may have been different (qualitative and quantitative) had the explants been sampled at 30 or 90 days, instead of 60 days. This timing of sampling effect was also observed in papaya by Heringer et al. (2013).

PGRs are an important determinant of the success of papaya tissue culture. Except for seed germination, which can take place on PGR-free basal medium, most organogenic experiments in papaya have employed PGRs. This is not an exhaustive exploration of the literature, but serves to highlight some of the diversity of PGRs explored to date. Phloridzin (a dihydrochalcone) at 3 g/L improved somatic embryo germination in ‘Maradol’ (Ascencio-Cabral et al. 2008). Auxins can induce rooting in *in vitro* explants while phloroglucinol, an auxin-like substance (Teixeira da Silva et al. 2013), improved rooting of established *in vitro* papaya plants (Teixeira da Silva 2013). An overview of the papaya literature, primarily from 2000 onwards, reveals that 2,4-D is the most commonly used PGR for inducing somatic embryogenesis, but the requirement for light and darkness appears to be not only genotype-dependent, but also dependent on other factors such as medium constituents and explant. In fact, very early studies such as Litz and Conover (1981, 1982, 1983), Fitch and Manshardt (1990) or Fitch (1993) already indicated the importance of 2,4-D and CW for somatic embryogenesis




Figure 1. The response of papaya (*Carica papaya* L.) 'Sunrise Solo' to single doses of select plant growth regulators. A. Seed germination. B. Leaf explant. C. Fasciated and/or hyperhydric shoots in response to 8 mg/L TDZ. D. SFC in response to 1 mg/L picloram. E. HC formation in response to 1 mg/L dicamba. F. HC formation in response to 2 mg/L GA₃. G. HC formation in response to 2 mg/L chitosan. H. HC formation in response to 4 mg/L 2,4-D. I. Root formation in response to 8% v/v CW. J. Leaves of developed adult plantlet: left = TDZ-derived; right = control. A-D = light; E, F = darkness. Leaf explants = G, H, I; stem tTCLs = C, D, E, F. HC = hard callus; SFC = soft, friable callus.

while Chen et al. (1987) noted that a combination of 1.0 mg/L NAA, 0.5 mg/L Kin and 1.0 mg/L GA₃ was essential for embryogenic callus induction. Khatoon and Sultana (1994) found that petiole segments and epidermal or longitudinal TCLs of 'Malir' could form callus and shoots in the presence of 2.5 mg/L NAA + 0.5 mg/L 2iP, and 1 mg/L NAA + 1 or 5 mg/L 2iP, respectively, all media requiring 15% (v/v) CW. Cabrera-Ponce et al. (1996) could propagate transgenic plantlets in the presence of 0.2 mg/L BA and 0.1 mg/L Kin. Fernando et al. (2001) found that 'Sunrise Solo' mature zygotic embryos could form somatic

embryos in the presence of 2 mg/L 2,4-D. Usman et al. (2002) claimed that leaves with mid-rib were much more receptive to somatic embryogenesis than petioles or leaves without mid-rib, as was also found in this study. Shoots of 'Honey Dew', 'Washington' and 'Co2' could be induced either with 2.25 μM TDZ or 4.4 μM BA + 0.5 μM NAA, elongated with 4.7 μM GA₃ and rooted with 14.7 μM IBA (Bhattacharya et al. 2003). Farzana et al. (2008) induced embryogenic callus in the presence of 3 mg/L NAA when 'Rathna' zygotic embryos and hypocotyls were used while 1 mg/L 2,4-D was optimal for young leaf explants. In that

study, NAA at 0.02 mg/L and 0.5 mg/L BA were best for somatic embryo maturation. Jattana et al. (2008) found that CW was important in the induction of embryogenic callus of 'Kaekdum' cotyledons, while somatic embryos could form in the presence of 2 mg/L 2,4-D and 4 mg/L Zea. Renukdas et al. (2010) found that several PGRs could be ranked in terms of their ability to form somatic embryos: 4 μ M picloram > 20 μ M dicamba > 10 μ M 2,4-D > 4 μ M 2,4,5-T; moreover, somatic embryos could mature in the presence of 1 μ M spermidine (a polyamine), 0.1 μ M ABA or 0.5 μ M AgNO₃ (an ethylene inhibitor). Malabadi et al. (2011) claimed successful somatic embryogenesis for 11 papaya cultivars in the presence of 4.52 μ M 2,4-D and 2.27 μ M TDZ. Mahadevamma et al. (2012) could induce friable callus from gamma ray-irradiated shoot tips using 6 mg/L BA and 2 mg/L IAA or 0.1 mg/L NAA and 0.6 mg/L BA. Wu et al. (2012) used 0.25 mg/L BA + 1 mg/L GA₃ to elongate shoots from somatic embryos of hermaphroditic 'Meizhonghong'. Heringer et al. (2013) claimed that polyethylene glycol is important for the maturation of hybrid UENF/CALIMAN 01 somatic embryos.

The response of papaya *in vitro* leaf tissues to a wide range of PGRs was tested. Several of these PGRs have not yet been used in papaya tissue culture, in part because of the wide success of somatic embryogenesis in response to 2,4-D, or shoot production in the presence of BA, with or without NAA. Papaya leaves respond sensitively to inductive agents such as PGRs *in vitro*, and thus serves as a useful model tropical fruit species. The ability to chemically control organogenesis, such as rhizogenesis (Teixeira da Silva 2013; Fig 11), or the responsiveness of papaya tissue cultures to LEDs and variation in the light spectrum (Teixeira da Silva JA 2014a) provide exciting prospects for papaya biotechnology. Reliable and flexible (i.e., alternative pathways) to achieve the same organogenic outcome serve well for genetic transformation experiments or studies on development, including those using TCLs (Teixeira da Silva et al. 2015).

ACKNOWLEDGEMENTS

The author thanks Prof. Michio Tanaka (Kagawa University, Japan) for research support. The author declares no conflicts of interest (financial or other).

REFERENCES

- Anandan R, Sudhakar D, Balasubramanian P, Gutiérrez-Mora A. 2012. *In vitro* somatic embryogenesis from suspension cultures of *Carica papaya* L. *Sci Hort* 136: 43-49.
- Ascencio-Cabral A, Gutiérrez-Pulido H, Rodríguez-Garay B, Gutiérrez-Mora A. 2008. Plant regeneration of *Carica papaya* L. through somatic embryogenesis in response to light quality, gelling agent and phloridzin. *Sci Hort* 118 (2): 155-160.
- Bhattacharya J, Renukdas NN, Khuspe SS, Rawal SK. 2003. Multiple shoot regeneration from immature embryo explants of papaya. *Biol Plant* 47 (3): 327-331.
- Cabrera-Ponce JL, Vegas-García A, Herrera-Estrella L. 1996. Regeneration of transgenic papaya plants via somatic embryogenesis induced by *Agrobacterium rhizogenes*. *In Vitro Cell Dev Biol Plant* 32: 86-90.
- Canini A, Alesiani D, D'Arcangelo G, Tagliatesta P. 2007. Gas chromatography-mass spectrometry analysis of phenolic compounds from *Carica papaya* L. leaf. *J Food Compos Anal* 20 (7): 584-590.
- Cha-um S, Kirdmanee C. 2007. Minimum growth *in vitro* culture for preservation of plant species. *Fruit Veg Cereal Sci Biotechnol* 1 (1): 13-25.
- Chen MH, Wang PJ, Maeda E. 1987. Somatic embryogenesis and plant regeneration in *Carica papaya* L. tissue culture derived from root explants. *Plant Cell Rep* 6: 348-351.
- de Almeida EP, de Oliveira RP, Dantas JLL. 2001. Papaya callus and somatic embryo induction and development. *Sci Agric* 58 (1): 51-54.
- Engelmann F. 2011. Use of biotechnologies for the conservation of plant biodiversity. *In Vitro Cell Dev Biol Pl* 47 (1): 5-16.
- Farzana ARF, Palkadapala PGVN, Meddegoda KMMN, Samarajeewa PK, Eeswara JP. 2008. Somatic embryogenesis in papaya (*Carica papaya* L. cv. Rathna). *J Nat Sci Found Sri Lanka* 36 (1): 41-50.
- Fernando JA, Melo M, Soares MKM, Appezzato-da-Glória B. 2001. Anatomy of somatic embryogenesis in *Carica papaya* L. *Braz Arch Biol Technol* 44 (3): 247-255.
- Fitch MMM. 1993. High frequency somatic embryogenesis and plant regeneration from papaya hypocotyl callus. *Plant Cell Tiss Org Cult* 32 (2): 205-212.
- Fitch MMM, Manshardt RM. 1990. Somatic embryogenesis and plant regeneration from immature zygotic embryos of papaya (*Carica papaya* L.). *Plant Cell Rep* 9 (6): 320-324.
- Giang DTT, Van PT, Tanaka M, Teixeira da Silva JA. 2011. Sterilization and germination of papaya (*Carica papaya* L.) seed and response to LEDs. *Seed Sci Biotechnol* 5 (1): 56-58.
- Heringer AS, Vale EM, Baroso T, Santa-Catarina C, Silveira V. 2013. Polyethylene glycol effects on somatic embryogenesis of papaya hybrid UENF/CALIMAN 01 seeds. *Theor Exp Plant Physiol* 25 (2): 116-124.
- Jattana A, Kijwijian B, Boadhipadma K, Leung DMW. 2008. Indirect somatic embryogenesis and synthetic seed production from cotyledon explants of papaya (*Carica papaya*) 'Kaekdum'. *J Appl Sci* 7 (2): 116-122.
- Jiménez VM, Mora-Newcomer E, Gutiérrez-Soto MV. 2014. Biology of the papaya plant. In: Ming R, Moore PH (eds), *Genetics and Genomics of Papaya*, Plant Genetics and Genomics: Crops and Models 10, Springer, New York.
- Khatoun K, Sultana R. 1994. Plant regeneration from *Carica papaya* cv Malir grown in tissue culture. *Pakistan J Bot* 26 (1): 191-195.
- Litz RE, Conover RA. 1981. Effect of sex type, season and other factors on *in vitro* establishment and culture of *Carica papaya* L. explants. *J Amer Soc Hort Sci* 106: 792-794.
- Litz RE, Conover RA. 1982. *In vitro* somatic embryogenesis and plant regeneration from *Carica papaya* L. ovular callus. *Plant Sci Lett* 26: 153-158.
- Litz RE, Conover RA. 1983. High frequency somatic embryogenesis from *Carica* suspension cultures. *Ann Bot* 51: 683-686.
- Mahadevamma M, Sahijram L, Kumari V, Shankarappa TH. 2012. *In vitro* mutation studies in papaya (*Carica papaya* L.). *CIB Tech J Biotechnol* 1 (1): 49-55
- Malabadi RB, Vijaykumar S, Mulgund GS, Nataraja K. 2011. Induction of somatic embryogenesis in papaya (*Carica papaya*). *Res Biotechnol* 2 (5): 40-55.
- Murashige T, Skoog F. 1962. A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiol Plant* 15: 473-499.
- Renukdas NN, Mohan ML, Khuspe SS, Rawal SK. 2010. Influence of phytohormones, culture conditions and ethylene antagonists on somatic embryo maturation and plant regeneration in papaya. *Intl J Agric Res* 5 (7): 511-520.
- Sharma S, Shahzad A, Teixeira da Silva JA. 2013. Synseed technology - A complete synthesis. *Biotech. Adv* 31: 186-207.
- Teixeira da Silva JA. 2012. Is BA (6-benzyladenine) BAP (6-benzylaminopurine)? *Asian Austr J Plant Sci Biotechnol* 6 (Special Issue 1): 121-124.
- Teixeira da Silva JA. 2013. *In vitro* rhizogenesis in papaya (*Carica papaya* L.). *J Plant Dev* 20: 51-55.
- Teixeira da Silva JA. 2014a. Photoauto-, photohetero- and photomixotrophic *in vitro* propagation of papaya (*Carica papaya* L.) and response of seed and seedlings to light-emitting diodes. *Thammasat Intl J Sci Technol* 19 (1): 57-71.
- Teixeira da Silva JA. 2014b. Callus induction in papaya (*Carica papaya* L.), and synseed production for low temperature storage and cryopreservation. *Folia Hort* 26 (2): 155-162.

- Teixeira da Silva JA, Altamura MM, Dobránszki J. 2015. The untapped potential of plant thin cell layers. *J Hort Res* 23 (2): 127-131.
- Teixeira da Silva JA, Dobránszki J. 2013a. Plant thin cell layers: a 40-year celebration. *J Plant Growth Reg* 32 (4): 922-943.
- Teixeira da Silva JA, Dobránszki J. 2013b. How timing of sampling can affect the outcome of the quantitative assessment of plant organogenesis. *Sci Hort* 159: 59-66.
- Teixeira da Silva JA, Dobránszki J. 2014. Dissecting the concept of the thin cell layer: theoretical basis and practical application of the plant growth correction factor to apple, *Cymbidium* and chrysanthemum. *J Plant Growth Reg* 33: 881-895.
- Teixeira da Silva JA, Dobránszki J. 2015. Plant thin cell layers: update and perspectives. *Folia Hort* 27 (2): 183-190.
- Teixeira da Silva JA, Dobránszki J, Ross S. 2013. Phloroglucinol in plant tissue culture. *In Vitro Cell Dev Biol Plant* 49 (1): 1-16.
- Teixeira da Silva JA, Rashid Z, Nhut DT, Sivakumar D, Gera A, Souza Jr. MT, Tennant PF. 2007. Papaya (*Carica papaya* L.) biology and biotechnology. *Tree For Sci Biotechnol* 1 (1): 47-73.
- Tennant PF. 2010. Transgenic papaya. *Transgenic Plant J* 4 (special issue 1): 1-96.
- Usman M, Fatima B, Jaskani MJ, Khan MM. 2002. Development of a callogenic protocol in papaya (*Carica papaya* L.). 1. *In vivo* grown vegetative explants. *Int J Agric Biol* 4 (1): 95-98.
- Wu K-L, Zeng S-J, Chen Z-L, Duan J. 2012. In vitro mass propagation of hermaphroditic *Carica papaya* cv. 'Meizhonghong'. *Pakistan J Bot* 44 (5): 1669-1676.