

Relationship between insect herbivory and environmental variables in forests of northern Iran

GOODARZ HAJIZADEH¹, HAMID JALILVAND¹, MOHAMMAD REZA KAVOSI²,
HASSAN BARIMANI VARANDI³

¹Department of Forestry, Faculty of Natural Resources, Sari Agricultural Sciences & Natural Resources University, P.O. Box:#578, Sari, Mazandaran Province, Iran. Tel. /Fax. +98 151 3822715 email: goodarzhajizadeh@gmail.com

²Department of Silviculture and Forest Ecology, Faculty of Forest Science, Gorgan Agricultural Sciences and Natural Resources University, Gorgan, Golestan Province, Iran.

³Agricultural and Natural Resources Research Center of Mazandaran, Sari, Mazandaran Province, Iran

Manuscript received: 28 May 2016. Revision accepted: 11 July 2016.

Abstract. Hajizadeh G, Jalilvand H, Kavosi MR, Varandi HB. 2016. Relationship between insect herbivory and environmental variables in forests of northern Iran. *Nusantara Bioscience* 8: 155-160. This research aimed to recognize the relationship between the intensity of insect herbivory and environmental variables (stand structure, and physical and chemical properties of soil and litter) in Hyrcanian forests, northern Iran. Three forest types (*Pinus brutia*, *Parrotia persica-Carpinus betulus*, and *Fagus orientalis*), which covered the elevational gradient of the study region, were selected. In each forest type, three random plots (20×20m) at 100m intervals were sampled. The dominant polyphagous species, *Ennomos quercinaria* (Hufnagel, 1767), belonging to Geometridae family was identified. The forest types experiencing the lowest and highest defoliation intensity were *P. brutia* (12±2.08) and *F. orientalis* (57.67±5.36), respectively. The intensity of insect herbivory was significantly correlated with various structural parameters [forest type, elevation, diameter at breast height (DBH), tree density, tree cutting and canopy opening (%)], physical and chemical properties of the soil [clay, pH, and potassium (K)], and litter properties [depth and total nitrogen (N)]. The study illustrates a way to monitor geometrid population dynamics across environmental gradients in terms of their population density, degree of polyphagy and distribution; such results could be useful for developing appropriate management plans for these economically important insects.

Keywords: Environmental gradient, forest type, Geometridae, outbreak, population dynamics

INTRODUCTION

The environmental condition is important to survival and abundance of insects. Environmental variables that affect population densities can be classified into three broad categories: (i) biotic; (ii) physical; and (iii) disturbances (Ciesla 2011). Biotic factors include host abundance and quality, levels of competition and natural enemies. Physical factors include climate, soil, and landscape features such as elevation, slope and aspect. Disturbances include wildfire, severe storms, and anthropogenic activities that can have a profound effect on the quality and quantity of food and breeding sites (Ciesla 2011).

In recent years, many studies have been conducted on the ecological and economic effects of insect defoliation on forest ecosystems. Outbreaks of defoliating insects are usually followed by a sudden reduction in tree growth and forest productivity (short-term effects), increased tree mortality or decline of tree health, changes in species composition and increased leaching of nutrients (long-term effects) (Swank et al. 1981; Webb et al. 1995; Eshleman et al. 1998; Reynolds et al. 2000). Herbivory levels can stimulate physiological responses in a range of host plants (Pataki et al. 1998; Chen et al. 2001; Ozaki et al. 2004a; Quentin et al. 2011). Allocation of carbon to various parts of the tree can be reduced (Schultz and Baldwin 1982)

because of restricted carbon storage above ground and increasing of soil microbial activity in below ground (Le Mellec et al. 2011). These effects on allocation and growth have economic importance. For example, in boreal forests in Canada, depletions caused by insect herbivory between 1977 and 1987 at an average rate of 106 m³/year had a substantial impact on timber in growth (Fleming 2000). In the USA, more than 20 million ha of forested land are annually affected by insect outbreaks, resulting in drastic economic losses of approximately US\$20 million per annum (Ayres and Lombardero 2000). In northeastern Germany, defoliation by the black arches moth (*Lymantria monacha*) estimated levels of defoliation of up to 500 000 ha, with economic losses of approximately €16 million (Kato 1993).

Several invasive insect species have caused extensive tree mortality in Hyrcanian forests, including *Lymantria dispar* (Hajizadeh et al. 2010), *Hyphantria cunea* (Kavosi and Gninenko 2007), *Altica viridula* (Kavosi 2007), and, more recently, species of Geometridae (*Ennomos quercinaria*, *Erannis defoliaria* and *Operophtera brumata*) (Barimani Varandi et al. 2006; Kiadaliri 2005, 2007; Rajaei et al. 2010; Babaei 2013). Most studies conducted in Hyrcanian forests have focused on identifying the insect pests and studying their biology and natural enemies. Therefore, the present study investigated whether there is a relationship between herbivory intensity and environmental

variables of stand structure, and the physical and chemical properties of soil and litter in Hyrcanian forests, northern Iran.

MATERIALS AND METHODS

Site study

The study area was located in the Darabkola forest, Sari, Northern Iran. The boundary of this area is located at 36°28′-36°33′ N and 53°16′-53°20′ W. The study region has an average annual temperature of 31°C, a total annual rainfall of 700-750 mm, and an altitudinal range of 140-880 m (asl.) (Figure 1). The Darabkola forest covers an area of approximately 2600 ha, comprising natural temperate and uneven aged stands. The main tree species are *Quercus castaneifolia*, *Carpinus betulus*, *Acer velutinum*, *Alnus subcordata*, *Tilia begonifolia*, *Parrotia persica*, *Ulmus glabra*, *Acer platanoides*, *Diospyros lotus*, *Zelkova carpinifolia*, *Fagus orientalis* and *Acer cappadocicum*.

Study design

A field survey carried out at Darabkola forest, Sari, northern Iran during the growing season, 2014 showed clear signs of herbivory and tree canopies that were almost devoid of leaves. To understand the relationship between herbivory intensity and environmental variables, three distinct forest types [*Pinus brutia* (150-400 m asl.), *Parrotia persica*-*Carpinus betulus* (450-600 m asl.), and

Fagus orientalis (650-900 m asl.)] were selected. In each forest type, three random plots (20×20m) were selected at 100m intervals and all trees within each plot were sampled. In addition, within each plot, environmental variables (stand structure, and physical and chemical soil and litter properties) were measured. Herbivory intensity based on Shariati Najafabadi et al. (2014) defined as follows: no defoliation (0), <10% (1), 11-25% (2), 26-40% (3), 41-60% (4), 61-80% (5) and >80% (6). For each tree, the diameter at breast height (DBH), tree height, density (number of stems), and tree cutting (to represent disturbance effects), were recorded. In the center of each plot, canopy opening cover (%) was measured. In each plot, three soil samples were taken from the depth of 0-10 cm and, after mixing, composed one soil sample (Maranon et al. 1999). Soil samples were air-dried and passed through a 2mm mesh. The following soil characteristics were determined (methods given in parentheses): (i) soil texture [hydrometric method (Bouyoucos 1962)]; (ii) soil pH (of a soil-water solution of 1:2.5); (iii) soil bulk density (BD) [clod method (Ghazanshahi 1999)]; (iv) total nitrogen (N) [semi-micro-Kjeldahl method (Bremner 1996)]; and (v) soil organic carbon (C) [Walkley and Black method (Walkley and Black 1934)]. In addition, available phosphorus (P) was analyzed according to standard methods (Olsen et al. 1954), and exchangeable potassium (K) was determined by ammonium acetate at pH 7.0 using a flame-photometer.




Figure 1. Study site in Mazandaran province, northern Iran. Sample plots in Darabkola forest are identified by red circles (Akbarimehr et al. 2016).

Data analysis

Normality of variables was checked using the Kolmogorov-Smirnov test, and Levene's test was used to examine the equality of variances. One-way analysis of variance (ANOVA) with a Tukey post-hoc test was used to examine differences in environmental variables between forest types. Correlations between insect herbivory and environmental variables were determined by Pearson's correlation coefficient. All statistical analyses were carried out in SPSS 19.0 (SPSS, Inc.) was used.

RESULTS AND DISCUSSION

Results

The dominant polyphagous species, *Ennomos quercinaria* (Hufnagel, 1767), belonging to Geometridae family was identified. In a comparison of stand structure characteristics, there was a significant difference in DBH and tree cutting ($p=0.05$) and density of trees ($p=0.01$) among the forest types. The lowest and highest of DBH were observed in *Pinus brutia* and *Parrotia persica-Carpinus betulus* forest types respectively. But, the maximum and minimum of tree cutting and tree density occurred in *Pinus brutia* and *Fagus orientalis* types respectively. Total N and depth of litter showed a

significant difference in forest types ($p=0.01$) (Table 1). The lowest and highest defoliation intensity were observed in *Pinus brutia* (12 ± 2.08) and *Fagus orientalis* (57.67 ± 5.36) types, respectively ($F=35.03$, $p=0.01$) (Table 2). No relation was found between herbivory intensity and height of trees, whereas significant correlations were found for other stand characteristics. Open stands ($p=0.05$) and moderate DBH classes ($p=0.01$) were attacked more by herbivores. In terms of the level of disturbance (represented by tree cutting), herbivorous populations were more abundant on trees with average levels of disturbance ($p=0.05$). Insect herbivory was increased with elevation (Figure 2). A significant correlation was found between herbivory intensity, clay soil, K ($p=0.01$), and pH ($p=0.05$). A significant relation was also found between litter depth and herbivory intensity ($p=0.01$) (Table 3).

Table 2. Comparison of defoliation intensity between three forest types

| Forest types (Elevation) | Mean \pm SE |
|-----------------------------------------------------------|-------------------|
| <i>Pinus brutia</i> (150-400 m asl.) | 12 \pm 2.08c |
| <i>Parrotia persica-Carpinus betulus</i> (400-650 m asl.) | 39.33 \pm 3.48b |
| <i>Fagus orientalis</i> (650-900 m asl.) | 57.67 \pm 5.36a |

Table 1. Comparison (mean \pm SE) of stand structure, physical and chemical soil and litter characteristics across forest types

| | Variable | <i>Pinus brutia</i> | <i>Parrotia persica-Carpinus betulus</i> | <i>Fagus orientalis</i> | F-value | P-value |
|-----------------|-------------------------|-----------------------------------|------------------------------------------|-------------------------------|----------------------|---------|
| Stand structure | Canopy opening (%) | 25 \pm 2.89 | 23.33 \pm 4.41 | 15 \pm 2.89 | 2.385 ^{ns} | 0.173 |
| | DBH (cm) | 13.15 \pm 0.7 ^b | 40.82 \pm 3.28 ^a | 39.29 \pm 7.71 ^a | 10.258 [*] | 0.012 |
| | Tree height (m) | 16.47 \pm 0.84 | 25.67 \pm 1.76 | 22.33 \pm 4.58 | 2.619 ^{ns} | 0.152 |
| | Density (tree/ha) | 1216.67 \pm 87.002 ^a | 316.67 \pm 58.33 ^b | 275 \pm 76.38 ^b | 50.533 ^{**} | 0.000 |
| | Tree cutting (ha) | 125 \pm 38.19 \pm 22.05 | 33.33 \pm 22.05 | 8.33 \pm 8.33 | 5.621 [*] | 0.042 |
| Soil | BD (g/cm ³) | 1.55 \pm 0.08 | 1.26 \pm 0.03 | 1.56 \pm 0.13 | 3.502 ^{ns} | 0.098 |
| | Clay (%) | 14.24 \pm 1.13 | 18.3 \pm 3.04 | 20.22 \pm 2.0003 | 1.9222 ^{ns} | 0.226 |
| | Silt (%) | 37.31 \pm 5.82 | 36.56 \pm 3.31 | 37.94 \pm 2.08 | 0.192 ^{ns} | 0.83 |
| | Sand (%) | 48.44 \pm 6.96 | 45.13 \pm 5.2 | 39.84 \pm 2.98 | 0.646 ^{ns} | 0.557 |
| | RH (%) | 24.51 \pm 0.79 | 29.15 \pm 6.78 | 25.34 \pm 2.61 | 0.243 ^{ns} | 0.792 |
| | pH (1:2.5) | 6.46 \pm 0.14 | 6.04 \pm 0.52 | 5.59 \pm 0.27 | 1.523 ^{ns} | 0.292 |
| | EC | 0.4 \pm 0.03 | 0.3 \pm 0.03 | 0.34 \pm 0.04 | 1.948 ^{ns} | 0.223 |
| | OC (%) | 5.97 \pm 1.15 | 5.94 \pm 0.64 | 4.29 \pm 0.39 | 1.461 ^{ns} | 0.304 |
| | N (%) | 0.1 \pm 0.8 | 0.03 \pm 0.01 | 0.04 \pm 0.01 | 1.59 ^{ns} | 0.279 |
| | OM | 10.28 \pm 1.98 | 10.24 \pm 1.11 | 7.39 \pm 0.67 | 1.461 ^{ns} | 0.304 |
| | C:N | 61.47 \pm 19.29 | 200.22 \pm 55.07 | 135.63 \pm 49.38 | 2.473 ^{ns} | 0.165 |
| | P (mg/kg) | 7.56 \pm 0.96 | 9.52 \pm 2.2 | 6.44 \pm 1.45 | 0.924 ^{ns} | 0.447 |
| K (mg/kg) | 204.72 | 144.52 \pm 33.57 | 124.86 \pm 26.48 | 2.67 ^{ns} | 0.148 | |
| Litter | RH (%) | 32 \pm 2.08 | 40.04 \pm 8.4 | 44.77 \pm 7.04 | 0.958 ^{ns} | 0.435 |
| | Depth (cm) | 0.97 \pm 0.15 ^b | 1.93 \pm 0.04 ^a | 2.63 \pm 0.32 ^a | 20.174 ^{**} | 0.002 |
| | OC (%) | 35.46 \pm 4.32 | 32.11 \pm 3.83 | 38.58 \pm 8.69 | 0.289 ^{ns} | 0.759 |
| | N (%) | 0.73 \pm 0.008 ^a | 0.48 \pm 0.008 ^c | 0.62 \pm 0.01 ^b | 104 ^{**} | 0.000 |
| | OM (%) | 61.13 \pm 7.45 | 55.35 \pm 6.61 | 66.51 \pm 14.99 | 0.289 ^{ns} | 0.759 |
| | C:N | 48.27 \pm 5.59 | 66.02 \pm 8.13 | 61.59 \pm 12.56 | 1.003 ^{ns} | 0.441 |
| | P (mg/kg) | 0.08 \pm 0.007 | 0.66 \pm 0.0006 | 0.06 \pm 0.009 | 3.414 ^{ns} | 0.102 |
| | K (mg/kg) | 0.2 | 0.01 \pm 0.005 | 0.01 \pm 0.005 | 2 ^{ns} | 0.216 |

Note: n.s. = not significant. * $p<0.05$, ** $p<0.01$. (ANOVA). BD (Bulk density), RH (Relative humidity), OC (Organic carbon), OM (Organic matter) and EC (Electrical conductivity)

Table 3. Pearson correlation coefficients between insect herbivory and stand structure, physical and chemical soil and litter characteristics across forest types

| | Characteristic | Pearson correlation | P value |
|-------------------|-------------------------|---------------------|---------------------|
| Stand structure | Forest type | 0.954 | 0.000** |
| | Elevation (asl.) | 0.934 | 0.000** |
| | Canopy opening (%) | -0.686 | 0.041* |
| | DBH (cm) | 0.798 | 0.01** |
| | Height (m) | 0.64 | 0.063 ^{ns} |
| | Density (tree/ha) | -0.886 | 0.001** |
| | Cutting tree (ha) | -0.705 | 0.034* |
| | Slope (%) | -0.392 | 0.296 ^{ns} |
| Soil properties | BD (g/cm ³) | -0.163 | 0.676 ^{ns} |
| | Clay (%) | 0.817 | 0.007 ^{ns} |
| | Silt (%) | 0.116 | 0.766 ^{ns} |
| | Sand (%) | -0.375 | 0.319 ^{ns} |
| | WHC (%) | 0.037 | 0.924 ^{ns} |
| | pH (1:2.5) | -0.683 | 0.042* |
| | EC | -0.422 | 0.256 ^{ns} |
| | C | -0.555 | 0.121 ^{ns} |
| | N (%) | -0.469 | 0.203 ^{ns} |
| | OM (%) | -0.555 | 0.121 ^{ns} |
| | C:N | 0.355 | 0.348 ^{ns} |
| | P (mg/kg) | -0.261 | 0.497 ^{ns} |
| K (mg/kg) | 0.822 | 0.007** | |
| Litter properties | Depth (cm) | 0.914 | 0.007** |
| | WHC | 0.328 | 0.398 ^{ns} |
| | C | 0.078 | 0.824 ^{ns} |
| | N (%) | -0.551 | 0.124 ^{ns} |
| | OM (%) | -0.078 | 0.842 ^{ns} |
| | C:N | 0.394 | 0.295 ^{ns} |
| | P (mg/kg) | -0.505 | 0.166 ^{ns} |
| | K (mg/kg) | -0.66 | 0.53 ^{ns} |

Note: ^{ns} not significant, *significant at $\alpha = 0.05$, **significant at $\alpha = 0.01$. BD (Bulk density), RH (Relative humidity), OC (Organic carbon), OM (Organic matter) and EC (Electrical conductivity)




Figure 2. The relationship between elevation and defoliation intensity

Discussion

The main purpose of this research was to elucidate the relationship between the intensity of phytophagous insect species and multiple biotic and abiotic variables in Hyrcanian forest. Our field survey showed the most of the herbivorous larvae collected belonged to the Geometridae moth's family.

In this study, between stand structures variables (DBH, elevation, trees density and height) and defoliation intensities the significant correlation was observed. Defoliation intensity was increased with elevation. This pattern probably related to microclimate, region conditions, disturbances, natural enemy, pure stand and especially host tree species. The primary host tree species of Geometridae family in the forest selected area was *Fagus orientalis* which covered the elevation of 650-900m (asl.). This result consistent with the other studies in northern Iran (Kavosi 2012). Insect population is known to reach a peak at mid-altitude gradients (Janzen et al. 1976; Holloway 1987; Olson 1994; Chey 2000). In addition, factors determining the insect herbivores at higher elevations are thought to include a reduction in host plant, reduced structural complexity of vegetation, climate conditions, and physiological adaptations. The possibility of lower predation at higher elevations might also be important in the abundances of the Geometridae in such regions. However, other studies have reported a reduction of insect at higher elevations. A significant relationship was also found between DBH and herbivory intensity, such that most of the defoliated trees were observed in moderate diameter classes. Karamiyan Omrani et al. (2014), reported that the relationship between environmental variables (diameter at breast height, trees height and elevation) and insect herbivory of Geometridae family in northern Iran, Gorgan province. Their results show that the highest defoliation occurred in *F. orientalis*, *C. betulus* and *P. persica* tree species. Also in this study, the most attacked trees were observed in moderate through high diameter classes. Positive relationship between defoliation intensity and environmental variables selected was observed. Moderate disturbance, unlike intensive disturbance, was followed by increasing of diversity and herbivory was consistent with the results of other studies (Sargeev 1998; Humphrey et al. 1999; Joshi et al. 1999; Kitahara and Sey 2001).

There are many mechanisms in which the activity of insect herbivores can cause changes in nutrient availability in soils: (i) insect herbivores can deposit significant quantities of grass into litter and soil (Fogal and Slansky 1985; Grace 1986). In the present study, N in litter and soil in *Fagus orientalis* plots was higher than in the other forest types. The relationship between insect herbivory and clay, pH, and K were also significantly correlated; (ii) insect herbivory can affect the nutrient content of rain that passes through the plant canopy (Tukey and Morgan 1963); (iii) insect herbivores may change the quantity and quality of litter fall from the canopy through defoliation, increase phenol content, root mortality, and changes in the relative abundance of plant species or genotypes with different

litter quality (Faeth et al. 1981); (iv) herbivory may influence plant populations not only by enhancing litter quality, but also through improving soil nutrient levels (Kielland et al. 1997); (v) herbivory affects relationship between plants and their symbionts (Bardgett et al. 1998; Hunter 2001); and (vi) herbivory affects tree canopy structure and ground covered as a result of changes in access to light and soil moisture. Changes in soil microclimate resulting from herbivory can affect the nutrient cycle. Similarly, by changing light availability, herbivory also affects the quality of litter by affecting plant chemistry (Hunter and Forkner 1999; Strand et al. 1999) and/or plant diversity and productivity (Van Der Wal et al. 2000). This hypothesis is more consistent with the results of the present study based on the significant correlation between herbivory intensity and canopy openness. In addition, our data do not show strong that changes in insect herbivory cause changes in nutrient cycling in soils. Which although a number of other published studies do support that conclusion.

In conclusion, elevation and tree density (stand structure), clay, K, pH (soil variables), and litter properties (C:N, depth, P, and K) are responsible for the observed defoliation in selected forest types. The study illustrates a way to monitor geometrid moth population dynamics across environmental gradients in terms of their population density, degree of polyphagy and distribution; such results could be useful for developing appropriate management plans for these economically important insects. In finally, variables that are correlated with insect herbivory can be measured to assess the relative risk of a forest for increased insect herbivory.

REFERENCES

- Akbarimehr M, Hosseini SA, Hojjati SM, Shariati F. 2016. Analysis of changes in stream water chemistry following forest management practices in Darabkola forest. *J Fac For Istanbul Univ* 66 (2): 539-547.
- Ayres MP, Lombardero MJ. 2000. Assessing the consequences of global change for forest disturbance from herbivores and pathogens. *Sci Total Environ* 262 (3): 263-286.
- Babaei MR. 2013. Complementary investigation on life cycle of *Ennomos quercinaria* (Lep. Geometridae) in Mazandaran province. *Iranian J For Range Prot Res* 11 (1): 1-9.
- Bardgett RD, Wardle DA, Yeates GW. 1998. Linking aboveground and below-ground interactions: how plant responses to foliar herbivory influence soil organisms. *Soil Biol Biochem* 30, 1867-1878.
- Barimani Varandi H, Babaei MR, Vatandoost A. 2006. Preliminary investigation on life cycle of *Ennomos quercinaria* (Lep: Geometridae) in Mazandaran. *Iranian J For Range Prot Res* 4 (2): 71-79.
- Bouyoucos CJ. 1962. Hydrometer method improved for making particle-size analysis of soil. *Agronomy* 54: 464-465.
- Bremner JM. 1996. Nitrogen-total. In: Sparks DL, Page AL, Helmke PA, Loeppert RH, Soltanpour PN, Tabatabai MA, Johnston CT, Sumner ME. (eds). *Methods of Soil Analysis*. Soil Science Society of America, Inc. American Society of Agronomy, Inc. Madison, Wisconsin, USA.
- Ciesla WM. 2011. *Forest Entomology: A Global Perspective*. Blackwell Publishing Ltd., Chichester, UK.
- Chen Z, Kolb TE, Clancy KM. 2001. Mechanisms of Douglas-fir resistance to western spruce budworm defoliation: bud burst phenology, photosynthetic compensation and growth rate. *Tree Physiol* 21: 1159-1169.
- Chey VK. 2000. Moth diversity in the tropical rain forest of Lanjak-Entimau, Sarawak, Malaysia. *Malayan Nat J* 54: 305-318.
- Eshleman KN, Morgan RP, Webb JR, Deviney FA, Galloway JN. 1998. Temporal patterns of nitrogen leakage from mid-Appalachian forested watersheds: role of insect defoliation. *Water Resour Res* 34: 2005-2116.
- Faeth SH, Connor EF, Simberloff DS. 1981. Early leaf abscission: a neglected source of mortality for folivores. *Amer Nat* 117: 409-415.
- Fleming RA. 2000. Climate change and insect disturbance regimes in Canada's boreal forests. *World Resour Rev* 12 (3): 521-548.
- Fogal WH, Slansky F. 1985. Contribution of feeding by European sawfly larvae to litter production and element flux on Scots pine plantations. *Canadian J For Res* 15: 484-487.
- Grace JR. 1986. The influence of gypsy moth on the composition and nutrient content of litter fall in a Pennsylvania oak forest. *For Sci* 32: 855-870.
- Ghazanshahi M. 1999. *Soil and Plant Analysis*. Motarjem, Tehran.
- Hajizadeh G. 2010. Comparison of integrated management and pheromone control on the intensity and spatial distribution of *Lymantria dispar* (L.) in Golestan province. [M.Sc. Thesis]. Gorgan Agricultural Science and Natural Science, Gorgan, Iran. [Persian].
- Holloway JD. 1987. Macrolepidoptera diversity in the Indo-Australian tropics: geographic, biotopic and taxonomic variations. *Biol J Linnean Soc* 30: 325-341.
- Humphrey JW, Hawes C, Peace AJ, Ferris KR, Jukes MR. 1999. Relationship between insect diversity and habitat characteristics in plantation forests. *Ecol Monogr* 113: 11-21.
- Hunter MD. 2001. Out of sight, out of mind: The impacts of root feeding insects in natural, managed systems. *Agricultural and Forest Entomology*, 3:3-10.
- Hunter MD, Forkner RE. 1999. Hurricane damage influences foliar polyphenolics and subsequent herbivory on surviving trees. *Ecology*, 80: 2676-2682.
- Janzen DH, Ataroff M, Farinas M, Reyes S, Rincon, N, Soler A, Soriano P, Vera M. 1976. Changes in the arthropod community along an elevational transect in the Venezuelan Andes. *Biotropica*, 8: 193-203.
- Joshi PC, Lockwood JA, Vashishth N, Singh A. 1999. Grasshopper (Orthoptera: Acridoidea) community dynamics in a most deciduous forest in India. *J Orthoptera Res* 8:17-23.
- Karamiyan Omrani L, Kavosi MR, Mohammadi J, Faridi F. 2014. Effects of responsible factors of Geometridae moth's intensity defoliation on four tree species in Ziarat forest, Gorgan, Northern Iran. The 1st National Conference on Stable Agriculture and Natural Resources. 30 January, Tehran. 8pp. [Persian].
- Kato F. 1993. Ansätze zur ökonomischen Bewertung der durch Schwammspinner und Nonne verursachten Schäden. *Mitteilungen aus der Biologischen Bundesanstalt fuer Land-und Forstwirtschaft Berlin-Dahlem (Germany)*, 293: 69-81.
- Kavosi MR. 2007. *Altica viridula* in Golestan Province. *Plant protection and Food*, 1 (1): 53.
- Kavosi MR, Gninenko YI. 2007. Hyphantria Cunea in Mazandaran and Ardabil Provinces. *Plant Protect Food* 1 (1): 52.
- Kavosi MR. 2012. Identification and control of the important forest pests and diseases in forest, nurses and national parks in northern Iran. Gorgan Agricultural Science and Natural Resources University press, Gorgan, Iran.
- Kiadaliri H, Ostovan H, Abaei M, Ahangaran Y. 2005. Investigation on the behaviour treat of leaf feeder moth (*Erannia defoliaria* clerck) and natural enemies in forests of the in west of Mazandaran province. *J agricultural Scis Islamic Azad University* 11 (1): 145-159.
- Kiadaliri H, Kazemi Najafi S, Ahangaran Y. 2007. The effect of leaf feeder moth (*Erannia defoliaria* & *Operophtera brumata*) on radial growth of three tree species in north of Iran (case study: Mashalakh, Nowshahr). *Iranian J For Poplar Res* 15 (3): 301-309.
- Kielland K, Bryant JP, Ruess RW. 1997. Moose herbivory and carbon turnover of early successional stands in interior Alaska. *Oikos* 80: 25-30.
- Kitahara M, Sei K. 2001. A comparison of the diversity and structure of butterfly communities in semi-natural and human-modified grassland habitats at the foot of Mt. Fuji, central Japan. *Biodiv Conserv* 10: 331-351.
- Le Mellec A, Gerold G, Michalzik B. 2011. Insect herbivory, organic matter deposition and effects on belowground organic matter fluxes in a central European oak forest. *Pl Soil* 342 (1-2): 393-403.

- Maranon T, Ajbilou R, Ojeda F, Arroyo J. 1999. Biodiversity of woody species in oak woodlands of southern Spain and northern Morocco. *Forest Ecology and Management* 115 (5): 147-156.
- Olsen SR, Cole CV, Watanabe FS, Dean LA. 1954. Estimation of available phosphorus in soil by extraction with sodium bicarbonate. U.S. Department of Agriculture Circular, Washington DC.
- Olson DM. 1994. The distribution of leaf litter invertebrates along a Neotropical altitudinal gradient. *J Trop Ecol* 10: 129-150.
- Ozaki K, Fukuyama K, Isono M, Takao G. 2004a. Simultaneous outbreak of three species web-spinning sawflies: influence of weather and stand structure. *For Ecol Manag* 187: 75-84.
- Pataki DE, Oren R, Phillips N. 1998. Responses of sap flux and stomatal conductance of *Pinus taeda* L. Trees to stepwise reductions in leaf area. *J Exp Bot* 49: 871-878.
- Quentin AG, O'Grady AP, Beadle CL, Worledge D, Pinkard EA. 2011. Responses of transpiration and canopy conductance to partial defoliation of *Eucalyptus globulus* trees. *Agric For Meteorol* 151: 356-364.
- Rajaei H, Abaai M, Hausmann A. 2010. First record of the winter moth *Operophtera brumata* (Linnaeus, 1758) (Lepidoptera: Geometridae) in northern Iran. *Iranian J Anim Biosyst* 6 (2): 61-66.
- Reynolds BC, Hunter MD, Crossley DA. 2000. Effects of canopy herbivory on nutrient cycling in a northern hardwood forest in western North Carolina. *Selbyana* 21: 74-78.
- Sargeev MG. 1998. Conservation of Orthoptera biological diversity relative to landscape change in temperate Eurasia. *J Insect Conserv* 2: 247-252.
- Schultz JC, Baldwin IT. 1982. Oak leaf quality declines in response to defoliation by gypsy moth larvae. *Science* 217 (4555): 149-151.
- Shariati Najafabadi H, Gorjestanizadeh S, Soltani A, Saeidi Z. 2014. Hawthorn preference of orchard ermine, *Yponomeuta padella* in Bâzoft forests. *Iranian J For Range Prot Res* 12 (1): 66-75.
- Strand M, Herms DA, Ayres MP, Kubiske ME, Kaufman MG, Walker ED, Pregitzer KS, Merritt RW. 1999. Effects of atmospheric CO₂ light availability and tree species on the quality of leaf detritus as a resource for tree hole mosquitoes. *Oikos* 84: 277-283.
- Swank WT, Waide JB, Crossley DA, Todd RL. 1981. Insect defoliation enhances nitrate export from forest ecosystems. *Oecologia* 51: 297-299.
- Tukey HB, Morgan JV. 1963. Injury to foliage and its effects upon the leaching of nutrients from above-ground plant parts. *Physiologia Planta* 16: 557-564.
- Van der Wal R, Egas M, Van der Veen A, Bakker JP. 2000. Effects of resource competition and herbivory on plant performance along a natural productivity gradient. *J Ecol* 88: 1-16.
- Walkley AJ, Black CA. 1934. Estimation of organic carbon by chromic acid titration method. *Soil Sci* 37 (22): 29-38.
- Webb JR, Cosby BJ, Deviney FA, Eshleman KN, Galloway JN. 1995. Change in the acid-base status of an Appalachian catchment following forest defoliation by the gypsy moth. *Water Air Soil Poll* 85: 535-540.